

ABOUT THE TEST FoundationOne®CDx is a next-generation sequencing (NGS) based assay that identifies genomic findings within hundreds of cancer-related genes.

PATIENT

DISEASE Prostate acinar adenocarcinoma

NAME

DATE OF BIRTH

SEX

MEDICAL RECORD #

PHYSICIAN

ORDERING PHYSICIAN

MEDICAL FACILITY

ADDITIONAL RECIPIENT

MEDICAL FACILITY ID

PATHOLOGIST

SPECIMEN

SPECIMEN SITE

SPECIMEN ID

SPECIMEN TYPE

DATE OF COLLECTION

SPECIMEN RECEIVED

Genomic Signatures

Microsatellite status - MS-Stable

Tumor Mutational Burden - 1 Muts/Mb

Gene Alterations

For a complete list of the genes assayed, please refer to the Appendix.

AR amplification

BRCA2 rearrangement intron 14

RET amplification

TMPRSS2 TMPRSS2-ERG fusion

MLH1 rearrangement intron 16

RAD21 amplification - equivocal†

TP53 L130R

† See About the Test in appendix for details.

12 Therapies approved in the EU

26 Clinical Trials

0 Therapies with Lack of Response

GENOMIC SIGNATURES

Microsatellite status - MS-Stable

Tumor Mutational Burden - 1 Muts/Mb

ACTIONABILITY

No therapies or clinical trials. see Genomic Signatures section

No therapies or clinical trials. see Genomic Signatures section

GENE ALTERATIONS	THERAPIES APPROVED IN THE EU (IN PATIENT'S TUMOR TYPE)	THERAPIES APPROVED IN THE EU (IN OTHER TUMOR TYPE)
AR - amplification	Abiraterone	none
	Bicalutamide	
	Cyproterone	
	Degarelix	
	Flutamide	
	Goserelin	
	Leuporelin	
	Triptorelin	
10 Trials see p. 12		
BRCA2 - rearrangement intron 14	none	Niraparib
		Olaparib
		Rucaparib
		Talazoparib
10 Trials see p. 15		
RET - amplification	none	none
3 Trials see p. 19		
TMPRSS2 - TMPRSS2-ERG fusion	none	none
10 Trials see p. 20		

GENE ALTERATIONS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIALS OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Alterations section.

MLH1 - rearrangement intron 16 p. 6 **TP53** - L130R p. 7
RAD21 - amplification - equivocal p. 7

NOTE Genomic alterations detected may be associated with activity of certain approved therapies; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and exhaustive. Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies.

Therapies contained in this report may have been approved through a centralized EU procedure or a national procedure in an EU Member State. Therapies, including but not limited to the following, have been approved nationally and may not be available in all EU Member States: Tretinoin, Anastrozole, Bicalutamide, Cyproterone, Exemestane, Flutamide, Goserelin, Letrozole, Leuporelin, Triptorelin.

PRF#

GENOMIC SIGNATURES

GENOMIC SIGNATURE

Microsatellite status

RESULT
MS-Stable

POTENTIAL TREATMENT STRATEGIES

On the basis of clinical evidence, MSS tumors are significantly less likely than MSI-H tumors to respond to anti-PD-1 immune checkpoint inhibitors¹⁻³, including approved therapies nivolumab and pembrolizumab⁴. In a retrospective analysis of 361 patients with solid tumors treated

with pembrolizumab, 3% were MSI-H and experienced a significantly higher ORR compared with non-MSI-H cases (70% vs. 12%, $p=0.001$)⁵.

FREQUENCY & PROGNOSIS

MSI has been reported in 3.1-14.6% of prostate cancer samples⁶⁻¹⁰. A study of prostate cancer in hereditary nonpolyposis colorectal cancer (HNPCC) families reported MSI-H in 4-50% of cases¹¹⁻¹³. The prognostic significance of MSI in prostate cancer has not been extensively studied (PubMed, May 2019).

FINDING SUMMARY

Microsatellite instability (MSI) is a condition of

genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor¹⁴. Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2¹⁴⁻¹⁶. This sample is microsatellite-stable (MSS), equivalent to the clinical definition of an MSS tumor: one with mutations in none of the tested microsatellite markers¹⁷⁻¹⁹. MSS status indicates MMR proficiency and typically correlates with intact expression of all MMR family proteins^{14,16,18-19}.

GENOMIC SIGNATURE

Tumor Mutational Burden

RESULT
1 Muts/Mb

POTENTIAL TREATMENT STRATEGIES

On the basis of strong clinical evidence, elevated TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-CTLA-4, anti-PD-L1, and anti-PD-1 immune checkpoint inhibitors; approved agents include ipilimumab, atezolizumab, avelumab, durvalumab, cemiplimab-rwlc, pembrolizumab, and nivolumab. Clinical studies have reported associations between elevated TMB and efficacy of PD-1- or PD-L1-targeting therapies, alone or in combination with other agents, in multiple types of solid tumors, including small cell²⁰⁻²¹ and non-small cell²²⁻³⁴ lung cancer, urothelial

carcinoma^{29,35-38}, melanoma^{25,29-30,39-43}, colorectal cancer^{29,44}, HNSCC^{29,45}, and other cancer types^{25,43,46-48}. For patients with melanoma, increased TMB has also been reported to be associated with clinical benefit from the CTLA-4 inhibitor ipilimumab^{32,49-51}.

FREQUENCY & PROGNOSIS

Prostate acinar adenocarcinoma harbors a median TMB of 2.7 mutations per megabase (mut/Mb), and 3.4% of cases have high TMB (>20 mut/Mb)⁵². Prostate cancer has been reported to harbor a relatively low TMB among solid tumors⁵³⁻⁵⁴, with approximately 0.5-1.5 (mut/Mb) in localized tumor samples⁵⁵⁻⁵⁷, and a higher but still low TMB of 2-5 mut/Mb in metastatic, castration-resistant prostate cancer (mCRPC) samples⁵⁸⁻⁶⁰. One study reported that 4 of 150 (2.7%) mCRPC cases harbored high TMB (nearly 50 mut/Mb), which was due to defects in mismatch repair genes MLH1 and MSH2 in 3 of the 4 cases⁶⁰. The effects of hypermutation on prognosis and clinical features in prostate cancer have not been extensively investigated (PubMed, Jul 2019).

FINDING SUMMARY

Tumor mutational burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma⁶¹⁻⁶² and cigarette smoke in lung cancer⁶³⁻⁶⁴, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes⁶⁵⁻⁶⁹, and microsatellite instability (MSI)^{65,68-69}. This sample harbors a low TMB. Compared to patients with tumors harboring higher TMB levels, patients with tumors harboring low TMB levels have experienced lower rates of clinical benefit from treatment with immune checkpoint inhibitors, including anti-CTLA-4 therapy in melanoma⁴⁹, anti-PD-L1 therapy in urothelial carcinoma³⁵, and anti-PD-1 therapy in non-small cell lung cancer and colorectal cancer^{4,64}.

PRF#

GENE ALTERATIONS

GENE

AR

ALTERATION

amplification

POTENTIAL TREATMENT STRATEGIES

Several approved drugs are available that target AR and/or the AR pathway in prostate cancer, including the antiandrogens apalutamide, bicalutamide, cyproterone, darolutamide, enzalutamide, flutamide, and nilutamide; the luteinizing hormone-releasing hormone (LHRH) agonists goserelin, leuprolide and triptorelin; the LHRH antagonist degarelix; and the CYP17A1 inhibitor abiraterone⁷⁰⁻⁷⁷. Resistance to androgen deprivation therapy (ADT) commonly occurs in prostate cancer through mechanisms such as increased AR expression, AR activation by tyrosine kinase-dependent signaling, alterations in AR co-activators, expression of alternatively spliced isoforms of AR mRNAs (AR-Vs), and extragonadal synthesis of androgenic compounds⁷⁸⁻⁸². AR signaling may promote radioresistance in prostate cancer by

transcriptionally upregulating DNA repair genes⁸³. There is preclinical evidence that development of resistance to anti-AR therapies, such as abiraterone and enzalutamide, may engender cross-resistance to the taxanes docetaxel and cabazitaxel⁸⁴; however, certain AR-Vs may remain sensitive to taxanes⁸⁵⁻⁸⁶. Approaches currently in clinical and preclinical development for prostate cancer include therapies that target AR nuclear translocation and degradation pathways⁸⁷⁻⁹³, combination approaches to suppress androgen biosynthesis⁹⁴, and the use of bromodomain and extraterminal (BET) inhibitors that disrupt the interaction between AR and BRD4⁹⁵⁻⁹⁷; the latter approach has potential to target AR-Vs⁹⁶⁻⁹⁸. The BET inhibitor mivebresib led to an SD rate of 60% (6/10) for patients with genomically unselected prostate cancer who had progressed on multiple standard of care therapies⁹⁹. Galeterone, a multifunctional AR inhibitor, has been evaluated in a number of Phase 1 and 2 studies and has been found to reduce prostate-specific antigen levels in 49-73% of patients, although clinical trials of this reagent are not recruiting patients¹⁰⁰.

FREQUENCY & PROGNOSIS

Aberrant activation of AR, through mutation and

amplification of AR, has been shown to be fundamental to prostate cancer progression; studies have reported AR gene amplification in 13-44% of patients with castration-resistant prostate cancer (CRPC)^{58,101-107}. AR amplification in prostate cancer appears to occur in response to androgen-deprivation therapy, as studies reported that 28-30% of patients with CRPC had AR amplification after, but not before, receiving androgen-deprivation therapy¹⁰²⁻¹⁰³. AR gene amplification is associated with advanced, hormone-refractory disease, and thus with poor prognosis. However, a study of prostate tumors that recurred during hormone therapy found no difference in patients with or without AR amplification in histological grade, Gleason score, or tumor stage¹⁰⁸.

FINDING SUMMARY

AR encodes the androgen receptor, a nuclear receptor that binds to testosterone and dihydroxytestosterone. AR is frequently amplified and overexpressed in castration-resistant prostate cancer (CRPC), also called hormone-refractory prostate cancer¹⁰⁹.

PRF#

GENE ALTERATIONS

GENE

BRCA2

ALTERATION

rearrangement intron 14

POTENTIAL TREATMENT STRATEGIES

Extensive clinical evidence in solid tumors indicates that inactivation of BRCA2 may confer sensitivity to PARP inhibitors, such as olaparib, rucaparib, or niraparib¹¹⁰⁻¹²⁶. Clinical response to PARP inhibitors has been reported for patients with either germline or somatic BRCA2 mutations^{111,116,119,126} and for patients who were platinum-resistant or refractory^{110,115,122,125}. Patients with BRCA-mutated platinum-resistant ovarian cancer have also responded to niraparib in combination with pembrolizumab; a Phase 1/2 study reported an ORR and a DCR of 45% (5/11) and 73% (8/11), respectively¹²⁷. In a Phase 2 trial of WEE1 inhibitor adavosertib in 31 patients with BRCA-mutated solid tumors, 1 patient with serous Fallopian tube cancer experienced a PR, and 4 patients (3 with ovarian cancer and 1 with solitary fibrous tumor) experienced SDs of greater than 6 months; patients with ovarian cancers had been previously treated with PARP inhibitors¹²⁸. Inactivation of BRCA2 may also predict sensitivity to DNA-damaging drugs such as the platinum chemotherapies cisplatin and carboplatin¹²⁹⁻¹³¹.

FREQUENCY & PROGNOSIS

BRCA2 mutations have been identified in 3–6% of primary and 6–7% of metastatic prostate cancer specimens^{57,60,132}, with deleterious germline BRCA2 mutations present in 5% of men with metastatic prostate cancer¹³³. The positive predictive value of prostate specific antigen (PSA) levels was found to be higher in patients with BRCA1/2 mutations than in the general population¹³⁴. BRCA2 germline mutations have been associated with attributes of aggressive prostate cancer at diagnosis, including high Gleason score, nodal involvement, advanced tumor stage, and metastatic spread¹³⁵. Germline BRCA2 mutation carriers had a significantly shorter cause-specific survival (CSS, 8.6 vs. 15.7 years) than noncarriers¹³⁵. Following radical conventional treatment for localized prostate cancer, patients with germline BRCA1/2 mutations experienced significantly shorter metastasis-free survival (HR=2.36) and CSS (HR=2.17) than noncarriers¹³⁶. For patients with metastatic castration-resistant prostate cancer (mCRPC), germline BRCA2 mutations were an independent marker of poor prognosis (CSS 17.4 vs. 33.2 months, HR=2.11) in 1 study¹³⁷. Germline BRCA2 mutations in mCRPC were associated with relative benefit from first-line abiraterone or enzalutamide compared with taxanes (CSS 24.0 vs. 17.0 months, PFS on the second systemic therapy 18.9 vs. 8.6 months) in a large prospective cohort study¹³⁷. Three patients with non-neuroendocrine prostate cancer harboring BRCA2 mutations derived clinical

benefit from treatment with platinum-based chemotherapy¹³⁸⁻¹³⁹.

FINDING SUMMARY

The BRCA2 tumor suppressor gene encodes a protein that regulates the response to DNA damage¹⁴⁰. Inactivating mutations in BRCA2 can lead to the inability to repair DNA damage and loss of cell cycle checkpoints, which can lead to tumorigenesis¹⁴¹. BRCA2 alterations that disrupt PALB2 binding (aa 21-39)¹⁴², the BRC repeats (aa 1002-2085), the DNA binding domain (aa 2479-3192), and/or the C-terminal RAD51 binding domain, as observed here, are predicted to be inactivating^{140,143-158}. Germline mutations in BRCA1 or BRCA2 are associated with breast-ovarian cancer familial susceptibility (BROVCA), also known as hereditary breast-ovarian cancer (HBOC)¹⁵⁹⁻¹⁶⁰. The lifetime risk of breast and ovarian cancer in BRCA1/2 mutation carriers has been estimated to be as high as 87% and 44%, respectively¹⁶¹, and elevated risk of other cancers, including gastric, pancreatic, prostate, and colorectal tumors, has been identified at frequencies of 20–60%¹⁶²⁻¹⁶⁹. The estimated prevalence of deleterious germline BRCA1/2 mutations in the general population is between 1:400 and 1:800, with an approximately 10-fold higher prevalence in the Ashkenazi Jewish population^{161,163,170-174}. In the appropriate clinical context, germline testing of BRCA2 is recommended.

GENE

RET

ALTERATION

amplification

POTENTIAL TREATMENT STRATEGIES

On the basis of clinical evidence, RET activating alterations may predict response to the multikinase inhibitors alectinib¹⁷⁵, cabozantinib¹⁷⁶⁻¹⁸³, lenvatinib¹⁸⁴, sorafenib¹⁸⁵, sunitinib¹⁸⁵⁻¹⁸⁶, and vandetanib¹⁸⁷⁻¹⁹³. Case reports

describe clinical benefit lasting 17 months with sunitinib for a patient with RET-amplified germ cell tumor¹⁹⁴ and stable disease from sunitinib and subsequent sorafenib for a patient with RET-amplified tongue adenocarcinoma¹⁸⁵.

FREQUENCY & PROGNOSIS

RET amplification and mutation have been reported in <1% of prostate carcinoma cases (cBioPortal, COSMIC, Oct 2019). Published data investigating the prognostic implications of RET alterations in prostate carcinoma are limited (PubMed, Dec 2018), although RET protein overexpression has been reported in both high-

grade prostate intraepithelial neoplasia (PIN) and in prostate carcinomas as compared to normal prostate tissue¹⁹⁵.

FINDING SUMMARY

RET (Rearranged during transfection) encodes a receptor tyrosine kinase primarily expressed in cells of the nervous system. It has been identified as a proto-oncogene that results in transformation of cells upon recombination with a partner gene¹⁹⁶. RET amplification has been reported to be amplified in cancer¹⁹⁷, and has been associated with responses to therapies that target RET^{185,194}.

PRF#

GENE ALTERATIONS

GENE

TMPRSS2

ALTERATION

TMPRSS2-ERG fusion

POTENTIAL TREATMENT STRATEGIES

There are no therapies to target rearrangement of TMPRSS2. Although direct targeting of ETS-family transcription factors is not possible, preclinical studies in prostate cancer with TMPRSS2-ETS family fusions have suggested sensitivity to agents targeting HDAC and PARP¹⁹⁸⁻²⁰⁰. Emerging clinical data indicate that 7 patients with ERG-rearranged prostate cancers achieved stable disease upon treatment with the PARP inhibitors niraparib or veliparib^{115,201}. Studies in prostate cancer have also reported that TMPRSS2-ETS family fusions may often co-occur with PTEN loss or PI3K pathway activation²⁰²⁻²⁰³.

HDAC and PARP inhibitors are in clinical trials in solid tumors. Limited preclinical data also suggest that bromodomain and extraterminal (BET) inhibitors suppress AR-BRD4 recruitment to the TMPRSS2 promoter, ERG expression, and AR-driven tumor growth in a preclinical castrate-resistant prostate cancer model⁹⁵.

FREQUENCY & PROGNOSIS

Fusions of TMPRSS2 with ETS family transcription factors are common in prostate carcinomas, reported in about 50-90% of cases²⁰⁴⁻²⁰⁷. TMPRSS2-ERG fusion is the most commonly observed fusion, reported in 50-59% of prostate cancers, but not in other tumor types examined^{204,208-213}. In addition, the TMPRSS2-ERG fusion has been identified in prostatic intraepithelial neoplasia but not in atypical adenomatous hyperplasia²¹⁴⁻²¹⁵. In general, a definitive relationship between TMPRSS2-ETS fusions and patient prognosis has been difficult to establish and is an area of ongoing study²¹⁶.

TMPRSS2-ERG has been associated with lower prostatic-specific antigen progression-free survival (PSA-PFS) in metastatic resistant prostate cancer patients treated with docetaxel or cabazitaxel²¹⁷.

FINDING SUMMARY

TMPRSS2 encodes an androgen-responsive serine protease that is most abundant in prostate tissue and normally expressed in other tissues such as colon, salivary gland, and stomach²¹⁸. It has been found to be downregulated in androgen-independent prostate cancer²¹⁹. The rearrangement seen here results in a fusion between TMPRSS2 and ERG, an ETS family transcription factor. TMPRSS2-ERG has been identified as a recurrent fusion in prostate cancer, in which the androgen-responsive promoter element of TMPRSS2 drives overexpression of ERG²⁰⁶. Different TMPRSS2-ERG fusions display varying degrees of oncogenicity²²⁰.

GENE

MLH1

ALTERATION

rearrangement intron 16

POTENTIAL TREATMENT STRATEGIES

MLH1 inactivation leads to MMR defects, high MSI, and increased mutational burden^{19,68,221-222}, which may predict response to the FDA-approved anti-PD-1 immunotherapies pembrolizumab and nivolumab^{4,64,223}. In a Phase 2 study of pembrolizumab in MSI-high colorectal cancer (CRC), three patients with MLH1 (germline) mutations experienced one partial response and two stable diseases⁴. Pembrolizumab demonstrated a significantly higher objective response rate in MSI-high CRC compared with microsatellite stable CRC (40% vs. 0%)⁴ and its efficacy correlated with high mutational burden in non-small cell lung cancer⁶⁴. Nivolumab achieved a complete response in a patient with MSI-high CRC²²³. Furthermore, MSI status correlates with higher PD-1 and PD-L1 expression¹, potential

biomarkers of response to anti-PD-1 immunotherapies. These therapies are in clinical trials for various tumor types and may be appropriate, particularly for hypermutant tumors.

FREQUENCY & PROGNOSIS

MLH1 mutation has been reported in 0-1.6% of prostate carcinoma cases^{55-56,58,60,107}. Approximately 12% (7/60) of prostate cancers were found to be hypermutated, with two of the hypermutated cases having MLH1 mutations⁸. MLH1 expression was shown to be reduced in approximately 50% of 39 prostate tumors analyzed in one study²²⁴. In another analysis, MLH1 nuclear expression was detected in 85% (6220/7275) prostate cancers; strong expression of MLH1 was observed in 55.2% of TMPRSS2-ERG-positive cases, but only in 32.9% of TMPRSS2-ERG-negative cases²²⁵. MLH1 expression was significantly associated with early biochemical recurrence in univariate analyses in one study²²⁵.

FINDING SUMMARY

MLH1 encodes the protein MutL homolog 1, colon cancer, nonpolyposis type 2, which binds PMS2 to form MutLalpha, a complex involved in DNA

mismatch repair (MMR)²²⁶. Defective MMR occurring as a result of mutation(s) in the MMR family (MLH1, MSH2, MSH6, or PMS2) can result in microsatellite instability (MSI), common in colon, endometrium, and stomach cancers¹⁵. MLH1 alterations that result in disruption or loss of the N-terminal ATPase-containing domain (amino acids 25-336)²²⁷⁻²²⁹, the exonuclease 1 (EXO1) interacting region or the C-terminal region necessary for PMS2 binding and formation of the MutL-alpha complex²³⁰, such as observed here, are predicted to be inactivating. Germline mutations in MLH1 are associated with a condition known as Lynch syndrome, which is characterized by increased risk of a number of cancers²³¹. Approximately 50% of Lynch syndrome-associated mutations have been attributed to alterations in MLH1²³². Lynch syndrome (also known as hereditary nonpolyposis colorectal cancer or HNPCC) accounts for 1-7% of all colorectal cancers and has an estimated prevalence in the general population between 1:600 and 1:2000²³³⁻²³⁵. In the appropriate clinical context, germline testing of MLH1 is recommended.

PRF#

GENE ALTERATIONS

GENE

RAD21

ALTERATION

amplification - equivocal

POTENTIAL TREATMENT STRATEGIES

There are no therapies to target alterations in this gene.

FREQUENCY & PROGNOSIS

RAD21 amplifications, point mutations, and truncating mutations have been reported in various cancers²³⁶. In the context of breast cancer, increased RAD21 expression has been correlated with poor prognosis in multiple subtypes²³⁷⁻²³⁸, including sporadic Grade 3 but not Grade 1 cancers²³⁷, as well as hereditary BRCA2-mutant and hereditary BRCA-wild-type but not hereditary BRCA1-mutant cancers²³⁷. Furthermore, SNPs in

or near RAD21 have been linked with risk of breast cancer development²³⁹⁻²⁴⁰. RAD21 overexpression has also been correlated with poor prognosis in endometrial cancer²⁴¹ and in colorectal cancer (CRC), especially in KRAS-mutant CRC²⁴². Heterogeneity of RAD21 expression also correlated with aggressive tumor behavior and shorter survival in endometrial cancer²⁴³. RAD21 amplification has been more frequently reported in hormone-refractory than in treatment-naïve prostate cancer, but RAD21 amplification did not correlate with expression²⁴⁴. In the context of ovarian cancer, both RAD21 overexpression and downregulation have been observed, but RAD21 expression was not prognostic²⁴⁵. Downregulation of RAD21 expression resulted in sensitization of cultured breast^{238,246} and CRC²⁴² cells to chemotherapy, thereby suggesting that RAD21 overexpression confers resistance to chemotherapy.

FINDING SUMMARY

RAD21 encodes a protein involved in DNA double-strand break repair and sister chromatid cohesion as a part of the cohesin complex²⁴⁷⁻²⁵⁰. In preclinical studies, downregulation of RAD21 or other cohesin components leads to loss of expression from amplified genes, as well as amplifications themselves upon cell passaging²⁵¹, but also leads to an increase in deletions, insertions, and other rearrangements²⁵². High RAD21 expression has also been associated with increased genomic instability²³⁷. Cohesin complex also organizes chromatin domains and regulates gene expression²⁵³⁻²⁵⁴. Both overexpression and reduction of expression of RAD21 has been reported to alter gene expression²⁵⁵. RAD21 amplification has been correlated with increased expression in breast^{237-238,256} and endometrial²⁴¹ cancers. Other RAD21 alterations, including truncating and point mutations, have been reported in the context of cancer, but the majority have not been characterized.

GENE

TP53

ALTERATION

L130R

TRANSCRIPT NUMBER

NM_000546

CODING SEQUENCE EFFECT

389T>G

12% (4/33) in patients who were TP53 wild-type²⁷¹. A Phase 2 trial of adavosertib in combination with chemotherapy (gemcitabine, carboplatin, paclitaxel, or doxorubicin) reported a 32% (30/94, 3 CR) ORR and a 73% (69/94) DCR in patients with platinum refractory TP53-mutated ovarian, Fallopian tube, or peritoneal cancer²⁷². A smaller Phase 2 trial of adavosertib in combination with carboplatin achieved a 43% (9/21, 1 CR) ORR and a 76% (16/21) DCR in patients with platinum-refractory TP53-mutated ovarian cancer²⁷³. The combination of adavosertib with paclitaxel and carboplatin in patients with TP53-mutated ovarian cancer also significantly increased PFS compared with paclitaxel and carboplatin alone²⁷⁴. A Phase 1 trial of neoadjuvant adavosertib in combination with cisplatin and docetaxel for head and neck squamous cell carcinoma (HNSCC) elicited a 71% (5/7) response rate in patients with TP53 alterations²⁷⁵. In a Phase 1b clinical trial of SGT-53 in combination with docetaxel in patients with solid tumors, 75% (9/12) of evaluable patients experienced clinical benefit, including 2 confirmed and 1 unconfirmed PRs and 2 instances of SD with significant tumor shrinkage²⁶⁵. Additionally, the combination of a CHK1 inhibitor and irinotecan reportedly reduced tumor growth and prolonged survival in a TP53-mutant, but not TP53-wild-type, breast cancer xenotransplant mouse model²⁷⁶.

FREQUENCY & PROGNOSIS

TP53 mutations have been reported in 18-40% of prostate cancers²⁷⁷⁻²⁷⁸. Overexpression of p53, which is indicative of TP53 dysregulation, has been reported to be significantly more common in late-stage and hormone-refractory prostate cancers and has been found to be associated with prostate-specific antigen (PSA) recurrence in low- and intermediate-grade prostate cancer²⁷⁹. TP53 loss has been found to be associated with prostate cancer-specific mortality in univariate analysis²⁸⁰.

FINDING SUMMARY

Functional loss of the tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers²⁸¹. Any alteration that results in the disruption or partial or complete loss of the region encoding the TP53 DNA-binding domain (DBD, aa 100-292) or the tetramerization domain (aa 325-356), such as observed here, is thought to dysregulate the transactivation of p53-dependent genes and is predicted to promote tumorigenesis²⁸²⁻²⁸⁴. Germline mutations in TP53 are associated with the very rare disorder Li-Fraumeni syndrome and the early onset of many cancers²⁸⁵⁻²⁸⁷, including sarcomas²⁸⁸⁻²⁹⁰. Estimates for the prevalence of germline TP53 mutations in the general population range from 1:5,000²⁹¹ to 1:20,000²⁹⁰. In the appropriate clinical context, germline testing of TP53 is recommended.

POTENTIAL TREATMENT STRATEGIES

There are no approved therapies to address TP53 mutation or loss. However, tumors with TP53 loss of function alterations may be sensitive to the WEE1 inhibitor adavosertib²⁵⁷⁻²⁶⁰, or p53 gene therapy and immunotherapeutics such as SGT-53²⁶¹⁻²⁶⁵ and ALT-801²⁶⁶. Missense mutations leading to TP53 inactivation may also be sensitive to therapies that reactivate mutant p53 such as APR-246²⁶⁷⁻²⁶⁹. In a Phase 1b trial in patients with p53-positive high-grade serous ovarian cancer, APR-246 combined with carboplatin and pegylated liposomal doxorubicin achieved a 52% (11/21) response rate and 100% DCR²⁷⁰. In a Phase 1 study, adavosertib in combination with gemcitabine, cisplatin, or carboplatin elicited PRs in 10% (17/176) and SDs in 53% (94/176) of patients with solid tumors; the response rate was 21% (4/19) in patients with TP53 mutations versus

PRF#

THERAPIES APPROVED IN THE EU

IN PATIENT'S TUMOR TYPE

Abiraterone

Assay findings association

AR
amplification

AREAS OF THERAPEUTIC USE

Abiraterone is an orally available CYP17 inhibitor. It is available in the EU in combination with prednisone or prednisolone to treat metastatic castration-resistant prostate cancer after progression on docetaxel-based chemotherapy, or after androgen blockade when chemotherapy is not indicated. It is also available in combination with prednisone or prednisolone and androgen deprivation therapy to treat newly diagnosed high-risk metastatic hormone-sensitive prostate cancer.

GENE ASSOCIATION

AR activation or amplification may predict sensitivity to abiraterone.

SUPPORTING DATA

The Phase 3 LATITUDE study in patients with previously untreated high-risk metastatic castration-naïve prostate cancer demonstrated that abiraterone plus prednisone and androgen deprivation therapy (ADT) significantly improved median OS (53.3 vs. 36.5 months, HR=0.66) and radiographic PFS (33.0 vs. 14.8 months, HR=0.47) compared to ADT plus placebo²⁹²⁻²⁹³. Phase 3 studies

have shown that abiraterone plus prednisone compared with placebo plus prednisone significantly improved median OS in patients with metastatic castration-resistant prostate cancer (mCRPC) who were either chemotherapy-naïve²⁹⁴ or had previously received chemotherapy²⁹⁵. For patients with non-metastatic CRPC, abiraterone plus prednisone also showed clinical efficacy in a Phase 2 study, with a median time to prostate-specific antigen (PSA) progression of 28.7 months and a median time to radiographic progressive disease of 41.4 months²⁹⁶. Abiraterone has also been evaluated to treat mCRPC in combination with other agents. A Phase 1b study evaluating abiraterone combined with docetaxel in chemotherapy-naïve mCRPC patients showed 86% (18/21) and 67% (14/21) of patients had $\geq 50\%$ and $\geq 90\%$ PSA decline from baseline, respectively²⁹⁷. In a Phase 1/2 study, abiraterone plus cabazitaxel showed further clinical efficacy (46% [12/26] PSA response rate and 21% [3/14] PR rate) in the post-docetaxel, post-abiraterone setting²⁹⁸. Treatment with cabazitaxel was more effective than abiraterone in patients with mCRPC that previously progressed on docetaxel and then enzalutamide (8.2 vs. 3.4 months, HR=0.44)²⁹⁹.

Bicalutamide

Assay findings association

AR
amplification

AREAS OF THERAPEUTIC USE

Bicalutamide is an orally available AR inhibitor that is available in the EU to treat advanced prostate cancer.

GENE ASSOCIATION

Clinical studies have demonstrated that AR activation or gene amplification may predict sensitivity to bicalutamide³⁰⁰⁻³⁰¹.

SUPPORTING DATA

A long-term follow-up to a Phase 3 clinical trial of bicalutamide/LHRH analog combination therapy in prostate carcinoma reported significant improvement in overall survival compared with LHRH analog monotherapy³⁰¹.

Cyproterone

Assay findings association

AR
amplification

AREAS OF THERAPEUTIC USE

Cyproterone is an orally available androgen receptor inhibitor that is available in the EU for the management of patients with prostate cancer.

GENE ASSOCIATION

AR activation or amplification may predict sensitivity to cyproterone.

SUPPORTING DATA

A Phase 2 study of cyproterone in combination with leuprolide for 36 weeks reported a five year survival rate of 80% in patients with prostate adenocarcinoma³⁰². A Phase 3 trial of 1045 patients with prostate cancer reported similar overall survival rates in patients treated with intermittent hormonal therapy, as compared to patients treated with cyproterone acetate³⁰³.

Degarelix

Assay findings association

AR
amplification

AREAS OF THERAPEUTIC USE

Degarelix is a gonadotrophin releasing hormone (GnRH) antagonist that is available in the EU to treat advanced hormone-dependent prostate cancer.

GENE ASSOCIATION

AR activation or amplification may predict sensitivity to

degarelix.

SUPPORTING DATA

A Phase 3 extension trial in patients with prostate cancer reported that degarelix was associated with significantly improved progression-free survival as compared with leuprolide³⁰⁴.

PRF#

THERAPIES APPROVED IN THE EU

IN PATIENT'S TUMOR TYPE

Flutamide

Assay findings association

AR
amplification

AREAS OF THERAPEUTIC USE

Flutamide is an orally available anti-androgen that is available in the EU to treat patients with advanced prostate cancer.

GENE ASSOCIATION

AR activation or amplification may predict sensitivity to flutamide.

SUPPORTING DATA

A Phase 2 clinical trial of the combination of suramin,

leuprolide, and flutamide in previously untreated patients with metastatic prostate cancer reported an overall response rate of 67%, include three complete responses and 30 partial responses³⁰⁵. A ten-year follow-up to a Phase 3 clinical trial in prostate carcinoma reported that long-term androgen-deprivation therapy (flutamide/goserelin) with radiation conferred significant benefit compared with short-term androgen-deprivation therapy with radiation³⁰⁶.

Goserelin

Assay findings association

AR
amplification

AREAS OF THERAPEUTIC USE

Goserelin is a luteinizing hormone-releasing hormone analog that is available in the EU to treat men with advanced prostate cancer and to treat premenopausal and perimenopausal women with hormone receptor-positive breast cancer.

GENE ASSOCIATION

AR activation or amplification may predict sensitivity to

goserelin.

SUPPORTING DATA

A ten-year follow-up to a Phase 3 clinical trial in prostate carcinoma reported that long-term androgen-deprivation therapy (flutamide/goserelin) with radiation conferred significant benefit compared with short-term androgen-deprivation therapy with radiation³⁰⁶.

Leuprorelin

Assay findings association

AR
amplification

AREAS OF THERAPEUTIC USE

Leuprorelin is an injectable luteinizing hormone-releasing hormone analog that is available in the EU to treat advanced hormone-dependent prostate cancer.

GENE ASSOCIATION

AR activation or amplification may predict sensitivity to leuprolide.

SUPPORTING DATA

A Phase 3 clinical trial reported that leuprolide administration was effective at lowering testosterone concentration to castrate levels in patients with prostate cancer³⁰⁷. A Phase 2 clinical trial of the combination of suramin, leuprolide, and flutamide in previously untreated patients with metastatic prostate cancer reported an overall response rate of 67%, include three complete responses and 30 partial responses³⁰⁵.

Triptorelin

Assay findings association

AR
amplification

AREAS OF THERAPEUTIC USE

Triptorelin is an injectable luteinizing hormone-releasing hormone analog that is available in the EU to treat advanced hormone-dependent prostate carcinoma.

GENE ASSOCIATION

AR activation or amplification may predict sensitivity to

triptorelin.

SUPPORTING DATA

A Phase 3 clinical trial reported that triptorelin administration was effective at lowering testosterone concentration to castration levels in patients with prostate cancer³⁰⁸.

PRF#

THERAPIES APPROVED IN THE EU IN OTHER TUMOR TYPE

Niraparib

Assay findings association

BRCA2

rearrangement intron 14

AREAS OF THERAPEUTIC USE

The PARP inhibitor niraparib is available in the EU for the maintenance treatment of patients with relapsed high grade serous epithelial ovarian, Fallopian tube, or primary peritoneal cancer who are in complete or partial response to platinum-based chemotherapy.

GENE ASSOCIATION

On the basis of clinical evidence in ovarian and breast cancers^{114-115,127}, loss or inactivation of either BRCA1 or BRCA2 may confer sensitivity to PARP inhibitors such as niraparib.

SUPPORTING DATA

In a Phase 2 study of niraparib in metastatic castration-

resistant prostate cancer (CRPC) that had progressed on at least 1 line of AR-targeted therapy in addition to at least 1 line of taxane chemotherapy, patients with biallelic BRCA1/2 mutation achieved a 62% (18/29) composite response rate, defined as ORR (38%, 6/16), circulating tumor cell (CTC) conversion (41%, 12/29), or a 50% decline in prostate-specific antigen level (PSA50) (52%, 15/29). Patients with non-BRCA biallelic DNA repair defects exhibited a 24% (5/21) composite RR, including a 13% (2/15) ORR, 19% (4/21) CTC conversion, and a 5% (1/21) PSA50³⁰⁹. In a Phase 1 study of niraparib treatment for patients with solid tumors, 43% (9/21) of patients with CRPC achieved SD, and 10 patients exhibited >30% decrease in CTCs¹¹⁵.

Olaparib

Assay findings association

BRCA2

rearrangement intron 14

AREAS OF THERAPEUTIC USE

The PARP inhibitor olaparib is available in the EU as maintenance therapy for patients with platinum-sensitive relapsed high-grade serous epithelial ovarian, Fallopian tube, or primary peritoneal cancer who are in complete or partial response to platinum-based chemotherapy, or as first-line maintenance for patients with these cancers who have a germline or somatic BRCA mutation and are in CR or PR after platinum-based chemotherapy. Olaparib is also approved to treat patients with HER2-negative advanced breast cancer and germline BRCA mutations who have been previously treated with chemotherapy; patients with hormone receptor-positive breast cancer should have been previously treated with, or considered not appropriate for, endocrine therapy.

GENE ASSOCIATION

Based on extensive clinical evidence in ovarian cancer¹²⁰⁻¹²⁴ as well as strong clinical evidence in multiple other cancer types^{110-112,120,123,310}, loss or inactivation of either BRCA1 or BRCA2 may confer sensitivity to olaparib.

SUPPORTING DATA

In pretreated patients with castrate-resistant prostate cancer (CRPC), a Phase 2 trial of olaparib reported PRs for 50% (4/8) and SDs for 25% (2/8) of heavily pretreated patients, with 88% (7/8) of these patients carrying germline BRCA1/2 mutation¹¹⁰. The TOPARP-B Phase 2 trial of olaparib reported a 46% (43/92) ORR and PFS of 5.4 months for pretreated patients with metastatic CRPC, and an 83% (25/30) ORR and PFS of 8.1 months for patients with CRPC and BRCA1/2 alterations³¹¹. Similarly,

another Phase 2 study of olaparib for biochemically recurrent prostate cancer observed a greater median prostate-specific antigen (PSA) PFS in men with BRCA2/ATM mutations (9 vs. 4 months without mutation)³¹². In another study, 33% (16/49) of patients with CRPC, including 88% (14/16) of patients with a mutation in 1 or more genes affecting homologous recombination benefited (PR, SD, and/or decline in PSA levels) from olaparib treatment¹¹¹. In a Phase 2 trial of olaparib combined with abiraterone in genomically unselected patients with CRPC, combination treatment improved median radiographic PFS (rPFS) compared to abiraterone and placebo (13.8 vs. 8.2 months, HR = 0.65); however, an increase in adverse events was noted in the combination group versus placebo³¹³. A Phase 1/2 trial combining olaparib with the immunotherapy durvalumab for patients with CRPC who were previously treated with abiraterone and/or enzalutamide reported a PSA decrease from a baseline of greater than or equal to 50% for 53% (9/17) of patients, including 4 RECIST PRs, and a median rPFS of 16.1 months; 3 responders harbored germline inactivating BRCA2 mutations with concurrent BRCA2 deletions of the second allele, while 2 responders displayed homozygous somatic BRCA2 inactivating alterations³¹⁴. A Phase 1 trial combined olaparib with the immunotherapy pembrolizumab for docetaxel-pretreated patients with metastatic CRPC and observed an ORR of 7% (2/28 PRs) and a PSA response rate of 14% (4/28 with 50% PSA decrease) in RECIST-evaluable patients, as well as a median rPFS of 4.7 months and median OS of 13.5 months in the overall population³¹⁵. Of note, homologous recombination deficiency was not detected in any of the patients³¹⁵.

PRF#

THERAPIES APPROVED IN THE EU

IN OTHER TUMOR TYPE

Rucaparib

Assay findings association

BRCA2

rearrangement intron 14

AREAS OF THERAPEUTIC USE

The PARP inhibitor rucaparib is available in the EU to treat patients with platinum-sensitive relapsed or progressive BRCA mutated (germline and/or somatic) high-grade epithelial ovarian, fallopian tube, or primary peritoneal cancer who have been treated with 2 or more prior lines of platinum-based chemotherapy and who are unable to tolerate further platinum-based chemotherapy. Rucaparib is also available for the maintenance treatment of patients with platinum sensitive relapsed high-grade epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in complete or partial response to platinum-based chemotherapy.

GENE ASSOCIATION

On the basis of strong clinical evidence in ovarian

cancer^{116-117,272}, as well as clinical data in other cancer types^{117,316-317}, loss or inactivation of either BRCA1 or BRCA2 may confer sensitivity to rucaparib.

SUPPORTING DATA

In the TRITON2 Phase 2 study for patients with metastatic castration-resistant prostate cancer (mCRPC) who previously progressed on androgen receptor-directed therapy and chemotherapy, rucaparib elicited an ORR of 28% (13/46) in an HRR-deficient population³¹⁸. Specifically, 44% (11/25) of patients with BRCA1/2 mutations, 1 patient with a FANCA mutation and 1 patient with a BRIP1 mutation experienced a confirmed RECIST response³¹⁸.

Talazoparib

Assay findings association

BRCA2

rearrangement intron 14

AREAS OF THERAPEUTIC USE

The PARP inhibitor talazoparib is available in the EU as monotherapy to treat patients with HER2-negative locally advanced or metastatic breast cancer with germline BRCA mutations, who have been previously treated with, or are not considered candidates for, available therapies.

GENE ASSOCIATION

On the basis of strong clinical data in breast cancer³¹⁹⁻³²¹ and additional clinical evidence in ovarian, pancreatic, and

prostate cancer³²²⁻³²⁴, loss or inactivation of either BRCA1 or BRCA2 may confer sensitivity to talazoparib.

SUPPORTING DATA

A study of talazoparib monotherapy reported 2 PRs in patients with BRCA-mutant prostate cancer patients³²². A case series in metastatic castration-resistant prostate cancer reported a decrease in the prostate-specific antigen (PSA) levels in one patient with a BRCA2 mutation after talazoparib treatment³²⁴.

NOTE Genomic alterations detected may be associated with activity of certain approved therapies; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Therapies listed in this report may not be complete and exhaustive and the therapeutic agents are not ranked in order of potential or predicted efficacy for this patient or in order of level of evidence for this patient's tumor type.

PRF#

CLINICAL TRIALS

IMPORTANT Clinical trials are ordered by gene and prioritized in the following descending order: Pediatric trial qualification → Geographical proximity → Trial phase → Trial verification within last 2 months. While every effort is made to ensure the accuracy of the information

contained below, the information available in the public domain is continually updated and should be investigated by the physician or research staff. The clinical trials listed in this report may not be complete and exhaustive or may include trials for which the patient does not meet the

clinical trial enrollment criteria. For additional information about listed clinical trials or to conduct a search for additional trials, please see clinicaltrials.gov or local registries in your region.

GENE

AR

ALTERATION

amplification

RATIONALE

Tumors with AR amplification or activation may be responsive to therapies that inhibit the

androgen receptor.

NCT03748641

PHASE 3

TARGETS
CYP17, PARP

A Study of Niraparib in Combination With Abiraterone Acetate and Prednisone Versus Abiraterone Acetate and Prednisone for Treatment of Participants With Metastatic Prostate Cancer

LOCATIONS: Calgary (Canada), Arizona, Vancouver (Canada), Victoria (Canada), California, Colorado, Connecticut, Florida, Kentucky, Maryland, Massachusetts, Nebraska, Nevada, New Jersey, New York, Hamilton (Canada), Kingston (Canada), Toronto (Canada), Pennsylvania, Montreal (Canada), South Carolina, Texas, Utah, Virginia, Washington, Buenos Aires (Argentina), Capital Federal (Argentina), Ciudad Autonomoma Buenos Aires (Argentina), Ciudad Autonomoma Buenos Aires (Argentina), Cordoba (Argentina), Mar del Plata (Argentina), Pergamino (Argentina), Rosario (Argentina), San Salvador de Jujuy (Argentina), Adelaide (Australia), Benowa (Australia), Birtinya (Australia), Brisbane (Australia), Darlinghurst (Australia), Hobart (Australia), Macquarie University (Australia), Malvern (Australia), Melbourne (Australia), Murdoch (Australia), Nedlands (Australia), Randwick (Australia), Wahroonga (Australia), Wollongong (Australia), Aalst (Belgium), Antwerpen (Belgium), Charleroi (Belgium), Gent (Belgium), Liege (Belgium), Merksem (Belgium), Natal (Brazil), Rio de Janeiro (Brazil), Sao Paulo (Brazil), Sorocaba (Brazil), São Paulo (Brazil), Pleven (Bulgaria), Plovdiv (Bulgaria), Sofia (Bulgaria), Vratsa (Bulgaria), Hradec Králove (Czechia), Liberec (Czechia), Olomouc (Czechia), Pardubice (Czechia), Plzen (Czechia), Praha 2 (Czechia), Praha 4 (Czechia), Besancon (France), Bordeaux (France), Clermont Ferrand (France), Lyon (France), Nancy (France), Nice Cedex 2 (France), Saint Herblain (France), Saint Mande (France), Strasbourg (France), Braunschweig (Germany), Duisburg (Germany), Magdeburg (Germany), Muenster (Germany), Nuertingen (Germany), Budapest (Hungary), Debrecen (Hungary), Gyula (Hungary), Nyiregyhaza (Hungary), Szeged (Hungary), Beer Yaakov (Israel), Haifa (Israel), Petah Tikva (Israel), Ramat Gan (Israel), Faenza (Italy), Milano (Italy), Napoli (Italy), Padova (Italy), Parma (Italy), Perugia (Italy), Pisa (Italy), Roma (Italy), Terni (Italy), Torino (Italy), Trento (Italy), Busan (Korea, Republic of), Daegu (Korea, Republic of), Daejeon (Korea, Republic of), Goyang-Si (Korea, Republic of), Gwangju (Korea, Republic of), Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Suwon (Korea, Republic of), George Town (Malaysia), Johor Bahru (Malaysia), Kota Kinabalu (Malaysia), Kuala Lumpur (Malaysia), Kuching (Malaysia), Aguascalientes (Mexico), Durango (Mexico), Monterrey (Mexico), Zapopan (Mexico), Amsterdam (Netherlands), Groningen (Netherlands), Leidschendam (Netherlands), Nieuwegein (Netherlands), Nijmegen (Netherlands), Rotterdam (Netherlands), Sittard-Geleen (Netherlands), Bydgoszcz (Poland), Gdańsk (Poland), Gdynia (Poland), Lodz (Poland), Siedlce (Poland), Szczecin (Poland), Warszawa (Poland), Wrocław (Poland), Barnaul (Russian Federation), Ivanovo (Russian Federation), Kursk (Russian Federation), Moscow (Russian Federation), Nizhny Novgorod (Russian Federation), Omsk (Russian Federation), Pyatigorsk (Russian Federation), Saint Petersburg (Russian Federation), Saint-Petersburg (Russian Federation), Saransk (Russian Federation), Sochi (Russian Federation), Tambov (Russian Federation), Tomsk (Russian Federation), Tyumen (Russian Federation), Vologda (Russian Federation), Johannesburg (South Africa), Pretoria (South Africa), Barcelona (Spain), Coruña (Spain), Jerez de la Frontera (Spain), Madrid (Spain), Málaga (Spain), Pozuelo de Alarcon (Spain), Sabadell (Spain), Santander (Spain), Valencia (Spain), Stockholm (Sweden), Uppsala (Sweden), Kaohsiung (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Taipei (Taiwan), Taoyuan (Taiwan), Adana (Turkey), Ankara (Turkey), Antalya (Turkey), Edirne (Turkey), Istanbul (Turkey), Izmir (Turkey), Kocaeli (Turkey), Cherkasy (Ukraine), Dnipro (Ukraine), Dnipro (Ukraine), Ivano-Frankivsk (Ukraine), Khakhiv (Ukraine), Kharkiv (Ukraine), Kyiv (Ukraine), Lviv (Ukraine), Poltava (Ukraine), Uzhgorod (Ukraine), Blackburn (United Kingdom), Lancaster (United Kingdom), London (United Kingdom), Torquay (United Kingdom), Truro (United Kingdom), Wolverhampton (United Kingdom)

PRF#

CLINICAL TRIALS
NCT03732820
PHASE 3

Study on Olaparib Plus Abiraterone as First-line Therapy in Men With Metastatic Castration-resistant Prostate Cancer

TARGETS
PARP, CYP17

LOCATIONS: Alabama, Alaska, Calgary (Canada), Edmonton (Canada), Arizona, Kelowna (Canada), California, Colorado, Connecticut, Illinois, Indiana, Louisiana, Michigan, Missouri, Montana, Nebraska, New Jersey, New York, North Carolina, Halifax (Canada), London (Canada), Toronto (Canada), Pennsylvania, Greenfield Park (Canada), Montreal (Canada), South Carolina, Texas, Wisconsin, Box Hill (Australia), Darlinghurst (Australia), Greenslopes (Australia), Herston (Australia), Kingswood (Australia), Kurralt Park (Australia), St Albans (Australia), Waratah (Australia), Gent (Belgium), Belo Horizonte (Brazil), Curitiba (Brazil), Fortaleza (Brazil), Porto Alegre (Brazil), Rio de Janeiro (Brazil), Sao Paulo (Brazil), São José do Rio Preto (Brazil), Santiago (Chile), Temuco (Chile), Viña del Mar (Chile), Brno (Czechia), Praha (Czechia), Praha 5 (Czechia), Angers Cedex 01 (France), BESANCON Cedex (France), Caen Cedex 05 (France), Marseille (France), Pierre Benite (France), Quimper Cedex (France), Toulouse Cedex 3 (France), Vandoeuvre les Nancy (France), Bergisch Gladbach (Germany), Bremen (Germany), Duisburg (Germany), Freiburg im Breisgau (Germany), Heinsberg (Germany), Köln (Germany), Mettmann (Germany), Nürnberg (Germany), Nürtingen (Germany), Ulm (Germany), Milano (Italy), Napoli (Italy), Orbassano (Italy), Pavia (Italy), Bunkyo-ku (Japan), Hirakata-shi (Japan), Kanazawa-shi (Japan), Kashiwara-shi (Japan), Kawagoe-shi (Japan), Kita-gun (Japan), Kyoto-shi (Japan), Maebashi-shi (Japan), Miyazaki-city (Japan), Nagoya-shi (Japan), Osaka-shi (Japan), Osakasayama-shi (Japan), Sagami-hara-shi (Japan), Sakura-shi (Japan), Shinjuku-ku (Japan), Toon-shi (Japan), Yokohama-shi (Japan), Daegu (Korea, Republic of), Goyang-si (Korea, Republic of), Seoul (Korea, Republic of), Hilversum (Netherlands), Nijmegen (Netherlands), Tilburg (Netherlands), Bratislava (Slovakia), Presov (Slovakia), Sala (Slovakia), Trencin (Slovakia), Barcelona (Spain), Gerona (Spain), Madrid (Spain), Malaga (Spain), Sevilla (Spain), Adana (Turkey), Ankara (Turkey), Istanbul (Turkey), Izmir (Turkey), Karsiyaka (Turkey), Guildford (United Kingdom), Manchester (United Kingdom), Sheffield (United Kingdom), Southampton (United Kingdom), Swansea (United Kingdom)

NCT03706365
PHASE 2

A Study of Abiraterone Acetate Plus Prednisone With or Without Abemaciclib (LY2835219) in Participants With Prostate Cancer

TARGETS
CYP17, CDK4, CDK6

LOCATIONS: Malaga (Spain), Arizona, Arkansas, Nürtingen (Germany), L'Hospitalet de Llobregat (Spain), California, Colorado, Connecticut, Plymouth (United Kingdom), Craiova (Romania), Florida, Nijmegen (Netherlands), Georgia, London (United Kingdom), Otopeni (Romania), Indiana, Kentucky, Seoul (Korea, Republic of), Chelsea (United Kingdom), Maryland, Massachusetts, Minnesota, Missouri, Nevada, Camperdown (Australia), Kogarah (Australia), Macquarie Park (Australia), Randwick (Australia), New York, Münster (Germany), North Carolina, Oklahoma, Oregon, Rotterdam (Netherlands), Tennessee, Texas, Utah, Vermont, Fitzroy (Australia), Copenhagen (Denmark), Næstved (Denmark), Tübingen (Germany), Utrecht (Netherlands), Bayamon (Puerto Rico), Bucuresti (Romania), Constanta (Romania), Barcelona (Spain), Madrid (Spain), Northampton (United Kingdom)

NCT00268476
PHASE 2/3

STAMPEDE: Systemic Therapy in Advancing or Metastatic Prostate Cancer: Evaluation of Drug Efficacy: A Multi-Stage Multi-Arm Randomised Controlled Trial

TARGETS
CYP17, AR

LOCATIONS: Reading (United Kingdom), Farnworth (United Kingdom), High Wycombe (United Kingdom), Cambridge (United Kingdom), Broomfield (United Kingdom), Chester (United Kingdom), Middlesbrough (United Kingdom), Carlisle (United Kingdom), Barnstaple (United Kingdom), Exeter (United Kingdom), Bournemouth (United Kingdom), Dorchester (United Kingdom), Poole (United Kingdom), Cottingham (United Kingdom), Eastbourne (United Kingdom), Saint Leonards-on-Sea (United Kingdom), Ashford (United Kingdom), Aylesbury (United Kingdom), Basingstoke (United Kingdom), Birmingham (United Kingdom), Brighton (United Kingdom), Burnley (United Kingdom), Burton-upon-Trent (United Kingdom), Bury St. Edmunds (United Kingdom), Crewe (United Kingdom), Darlington (United Kingdom), Derby (United Kingdom), Doncaster (United Kingdom), Dudley (United Kingdom), Gloucester (United Kingdom), Hereford (United Kingdom), Kidderminster (United Kingdom), Leeds (United Kingdom), Leicester (United Kingdom), Liverpool (United Kingdom), London (United Kingdom), Manchester (United Kingdom), Shrewsbury (United Kingdom), Stockport (United Kingdom), Sunderland (United Kingdom), Torquay (United Kingdom), Warrington (United Kingdom), Whitehaven (United Kingdom), Wigan (United Kingdom), Worcester (United Kingdom), Worthing (United Kingdom), Harlow (United Kingdom), Romford (United Kingdom), Westcliff-On-Sea (United Kingdom), Swansea (United Kingdom), Cheltenham (United Kingdom), Chur (Switzerland), Oldham (United Kingdom), Southampton (United Kingdom), Stevenage (United Kingdom), Inverness (United Kingdom), Newport (United Kingdom), Steeton (United Kingdom), Canterbury (United Kingdom), Maidstone (United Kingdom), Margate (United Kingdom), Glasgow (United Kingdom), Preston (United Kingdom), Southport (United Kingdom), Northwood (United Kingdom), Edinburgh (United Kingdom), Newcastle (United Kingdom), Scarborough (United Kingdom), Belfast (United Kingdom), Nottingham (United Kingdom), Sutton-in-Ashfield (United Kingdom), Oxford (United Kingdom), Cosham (United Kingdom), Ayr (United Kingdom), Bath (United Kingdom), Bristol (United Kingdom), Taunton (United Kingdom), Weston Super Mare (United Kingdom), Yeovil (United Kingdom), Sheffield (United Kingdom), Stoke-on-Trent (United Kingdom), Ipswich (United Kingdom), Guildford (United Kingdom), Sutton (United Kingdom), Newcastle-Upon-Tyne (United Kingdom), South Shields (United Kingdom), Lausanne (Switzerland), Aberystwyth (United Kingdom), Cardiff (United Kingdom), Sutton Coldfield (United Kingdom), Bradford (United Kingdom), Huddersfield (United Kingdom), Swindon (United Kingdom), Bebington (United Kingdom), Winterthur (Switzerland), Aarau (Switzerland), Basel (Switzerland), Berne (Switzerland), Liestal (Switzerland), St. Gallen (Switzerland), Zurich (Switzerland), Barnet (United Kingdom), Colchester (United Kingdom), Larbert (United Kingdom), Lincoln (United Kingdom), Stockton-on-Tees (United Kingdom), Wolverhampton (United Kingdom)

PRF#

CLINICAL TRIALS
NCT02799706
PHASE 3

Trial Comparing Irradiation Plus Long Term Adjuvant Androgen Deprivation With GnRH Antagonist Versus GnRH Agonist Plus Flare Protection in Patients With Very High Risk Localized or Locally Advanced Prostate Cancer

TARGETS
LHRH

LOCATIONS: Brussel (Belgium), Brussels (Belgium), Haine-Saint-Paul (Belgium), Kortrijk (Belgium), Namur (Belgium), Wilrijk (Belgium), Copenhagen (Denmark), Amiens (France), Arras (France), Bayonne (France), Beuvry (France), Grenoble (France), Lyon (France), Nantes (France), Paris (France), Saint-Gregoire (France), Toulouse (France), Berlin (Germany), Freiburg (Germany), Magdeburg (Germany), Firenze (Italy), Meldola (Italy), Alcorcón (Spain), Barcelona (Spain), Bilbao (Spain), Cartagena (Spain), Córdoba (Spain), Las Palmas De Gran Canaria (Spain), Pamplona (Spain), Sabadell (Spain), Salamanca (Spain), Terrassa (Spain), Vigo (Spain), Bellinzona (Switzerland), Bern (Switzerland), Geneve (Switzerland), Nottingham (United Kingdom)

NCT03568656
PHASE 1/2

Study to Evaluate CCS1477 in Advanced Tumours

TARGETS
CYP17, CBP, p300, AR

LOCATIONS: Belfast (United Kingdom), Leicester (United Kingdom), Newcastle (United Kingdom), Sutton (United Kingdom)

NCT03070886
PHASE 2/3

Antiandrogen Therapy and Radiation Therapy With or Without Docetaxel in Treating Patients With Prostate Cancer That Has Been Removed by Surgery

TARGETS
LHRH, AR

LOCATIONS: Alabama, Alaska, Arizona, Arkansas, California, Colorado, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Iowa, Kansas, Louisiana, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, South Carolina, South Dakota, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, San Juan (Puerto Rico)

NCT03009981
PHASE 3

A Study of Androgen Annihilation in High-Risk Biochemically Relapsed Prostate Cancer

TARGETS
LHRH, AR, CYP17

LOCATIONS: Arizona, California, District of Columbia, Hawaii, Illinois, Kansas, Maine, Massachusetts, Minnesota, Missouri, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Ohio, Oklahoma, Oregon, Pennsylvania, South Carolina, Texas, Wisconsin

NCT03414034
PHASE 2

Onvansertib in Combination With Abiraterone and Prednisone in Adult Patients With Metastatic Castration-Resistant Prostate Cancer

TARGETS
PLK1, CYP17

LOCATIONS: Massachusetts

NCT03436654
PHASE 2

Multi-arm Multi-modality Therapy for Very High Risk Localized and Low Volume Metastatic Prostatic Adenocarcinoma

TARGETS
AR, CYP17

LOCATIONS: Illinois, Massachusetts, New York, Ohio

PRF#

CLINICAL TRIALS
GENE
BRCA2
RATIONALE
BRCA2 loss or inactivating alterations may

predict sensitivity to PARP inhibitors.

ALTERATION
rearrangement intron 14

NCT03395197
PHASE 3

Talazoparib + Enzalutamide vs. Enzalutamide Monotherapy in mCRPC (TALAPRO-2)

TARGETS
PARP

LOCATIONS: Santiago de Compostela (Spain), Nagoya (Japan), Alaska, Hiroasaki (Japan), Arizona, Tauranga (New Zealand), L'Hospitalet de Llobregat (Spain), Sabadell (Spain), Pergamino (Argentina), Cremona (Italy), California, Christchurch (New Zealand), Kashiwa (Japan), Colorado, Port Elizabeth (South Africa), Meldola (Italy), Goyang-si (Korea, Republic of), Sapporo (Japan), Indiana, Kanazawa (Japan), Yokohama (Japan), Yokosuka (Japan), Kentucky, Nebraska, New Jersey, New Mexico, Camperdown (Australia), Darlinghurst (Australia), Port Macquarie (Australia), New York, Ohio, Osaka-shi (Japan), Osakasayama (Japan), Suita (Japan), Pennsylvania, Auchenflower (Australia), Brisbane (Australia), Chermiside (Australia), South Brisbane (Australia), Southport (Australia), Rosario (Argentina), Hamamatsu (Japan), South Carolina, Terni (Italy), Orbassano (Italy), Tennessee, Texas, Meguro-ku (Japan), Shinjuku-ku (Japan), Utah, Vina del Mar (Chile), Melbourne (Australia), North Melbourne (Australia), Hamilton (New Zealand), Washington, Cape Town (South Africa), George (South Africa), Kraaifontein, Cape Town (South Africa), Wisconsin, Cordoba (Argentina), Gent (Belgium), Kortrijk (Belgium), Yvoir (Belgium), Ostrava-Poruba (Czechia), Helsinki (Finland), Kempele (Finland), Kuopio (Finland), Oulu (Finland), Tampere (Finland), Turku (Finland), Suresnes (France), Suresnes Cedex (France), VILLEJUIF cedex (France), Kirchheim (Germany), Muenster (Germany), Nuertingen (Germany), Ravensburg (Germany), Reutlingen (Germany), Budapest (Hungary), Debrecen (Hungary), Pecs (Hungary), Tel Aviv (Israel), Napoli (Italy), Chiba (Japan), Fukuoka (Japan), Kagoshima (Japan), Kumamoto (Japan), Tokushima (Japan), Yamagata (Japan), Busan (Korea, Republic of), Daegu (Korea, Republic of), Seoul (Korea, Republic of), Auckland (New Zealand), Lorenskog (Norway), Norbyhagen (Norway), Oslo (Norway), Trondheim (Norway), Arequipa (Peru), Lima (Peru), Gdynia (Poland), Otwock (Poland), Warszawa (Poland), Barcelona (Spain), Solna (Sweden), Stockholm (Sweden), Umeå (Sweden), Cornwall (United Kingdom), Cronwall (United Kingdom), Glasgow (United Kingdom), London (United Kingdom)

NCT03748641
PHASE 3

A Study of Niraparib in Combination With Abiraterone Acetate and Prednisone Versus Abiraterone Acetate and Prednisone for Treatment of Participants With Metastatic Prostate Cancer

TARGETS
CYP17, PARP

LOCATIONS: Calgary (Canada), Arizona, Vancouver (Canada), Victoria (Canada), California, Colorado, Connecticut, Florida, Kentucky, Maryland, Massachusetts, Nebraska, Nevada, New Jersey, New York, Hamilton (Canada), Kingston (Canada), Toronto (Canada), Pennsylvania, Montreal (Canada), South Carolina, Texas, Utah, Virginia, Washington, Buenos Aires (Argentina), Capital Federal (Argentina), Ciudad Automoma Buenos Aires (Argentina), Ciudad Autonoma Buenos Aires (Argentina), Cordoba (Argentina), Mar del Plata (Argentina), Pergamino (Argentina), Rosario (Argentina), San Salvador de Jujuy (Argentina), Adelaide (Australia), Benowa (Australia), Birtinya (Australia), Brisbane (Australia), Darlinghurst (Australia), Hobart (Australia), Macquarie University (Australia), Malvern (Australia), Melbourne (Australia), Murdoch (Australia), Nedlands (Australia), Randwick (Australia), Wahroonga (Australia), Wollongong (Australia), Aalst (Belgium), Antwerpen (Belgium), Charleroi (Belgium), Gent (Belgium), Liege (Belgium), Merksem (Belgium), Natal (Brazil), Rio de Janeiro (Brazil), Sao Paulo (Brazil), Sorocaba (Brazil), São Paulo (Brazil), Pleven (Bulgaria), Plovdiv (Bulgaria), Sofia (Bulgaria), Vratsa (Bulgaria), Hradec Králove (Czechia), Liberec (Czechia), Olomouc (Czechia), Pardubice (Czechia), Plzen (Czechia), Praha 2 (Czechia), Praha 4 (Czechia), Besancon (France), Bordeaux (France), Clermont Ferrand (France), Lyon (France), Nancy (France), Nice Cedex 2 (France), Saint Herblain (France), Saint Mande (France), Strasbourg (France), Braunschweig (Germany), Duisburg (Germany), Magdeburg (Germany), Muenster (Germany), Nuertingen (Germany), Budapest (Hungary), Debrecen (Hungary), Gyula (Hungary), Nyiregyhaza (Hungary), Szeged (Hungary), Beer Yaakov (Israel), Haifa (Israel), Petah Tikva (Israel), Ramat Gan (Israel), Faenza (Italy), Milano (Italy), Napoli (Italy), Padova (Italy), Parma (Italy), Perugia (Italy), Pisa (Italy), Roma (Italy), Terni (Italy), Torino (Italy), Trento (Italy), Busan (Korea, Republic of), Daegu (Korea, Republic of), Daejeon (Korea, Republic of), Goyang-Si (Korea, Republic of), Gwangju (Korea, Republic of), Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Suwon (Korea, Republic of), George Town (Malaysia), Johor Bahru (Malaysia), Kota Kinabalu (Malaysia), Kuala Lumpur (Malaysia), Kuching (Malaysia), Aguascalientes (Mexico), Durango (Mexico), Monterrey (Mexico), Zapopan (Mexico), Amsterdam (Netherlands), Groningen (Netherlands), Leidschendam (Netherlands), Nieuwegein (Netherlands), Nijmegen (Netherlands), Rotterdam (Netherlands), Sittard-Geleen (Netherlands), Bydgoszcz (Poland), Gdańsk (Poland), Gdynia (Poland), Lodz (Poland), Siedlce (Poland), Szczecin (Poland), Warszawa (Poland), Wrocław (Poland), Barnaul (Russian Federation), Ivanovo (Russian Federation), Kursk (Russian Federation), Moscow (Russian Federation), Nizhny Novgorod (Russian Federation), Omsk (Russian Federation), Pyatigorsk (Russian Federation), Saint Petersburg (Russian Federation), Saint-Petersburg (Russian Federation), Saransk (Russian Federation), Sochi (Russian Federation), Tambov (Russian Federation), Tomsk (Russian Federation), Tyumen (Russian Federation), Vologda (Russian Federation), Johannesburg (South Africa), Pretoria (South Africa), Barcelona (Spain), Coruña (Spain), Jerez de la Frontera (Spain), Madrid (Spain), Málaga (Spain), Pozuelo de Alarcon (Spain), Sabadell (Spain), Santander (Spain), Valencia (Spain), Stockholm (Sweden), Uppsala (Sweden), Kaohsiung (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Taipei (Taiwan), Taoyuan (Taiwan), Adana (Turkey), Ankara (Turkey), Antalya (Turkey), Edirne (Turkey), Istanbul (Turkey), Izmir (Turkey), Kocaeli (Turkey), Cherkasy (Ukraine), Dnipro (Ukraine), Dnipro (Ukraine), Ivano-Frankivsk (Ukraine), Khakhiv (Ukraine), Kharkiv (Ukraine), Kyiv (Ukraine), Lviv (Ukraine), Poltava (Ukraine), Uzhgorod (Ukraine), Blackburn (United Kingdom), Lancaster (United Kingdom), London (United Kingdom), Torquay (United Kingdom), Truro (United Kingdom), Wolverhampton (United Kingdom)

PRF#

CLINICAL TRIALS
NCT02975934
PHASE 3

A Study of Rucaparib Verses Physician's Choice of Therapy in Patients With Metastatic Castration-resistant Prostate Cancer and Homologous Recombination Gene Deficiency

TARGETS
CYP17, PARP, AR

LOCATIONS: Alabama, Calgary (Canada), Arizona, California, Colorado, Connecticut, Delaware, London (United Kingdom), Oxford (United Kingdom), Preston (United Kingdom), Slough (United Kingdom), Florida, Hawaii, Louisiana, Winnipeg (Canada), Maryland, Massachusetts, Michigan, Minnesota, Nebraska, Nevada, Moncton (Canada), Miranda (Australia), Orange (Australia), Saint Leonards (Australia), New York, North Carolina, Ohio, London (Canada), Ottawa (Canada), Toronto (Canada), Oregon, Tennessee, Texas, Virginia, Washington, Subiaco (Australia), Frankston (Australia), Geelong (Australia), Hobart (Australia), Wagga Wagga (Australia), Antwerp (Belgium), Kortrijk (Belgium), Liège (Belgium), Namur (Belgium), Oshawa (Canada), Copenhagen (Denmark), Herlev (Denmark), Vejle (Denmark), Dijon (France), Le Mans (France), Lille (France), Lyon (France), Nancy (France), Paris (France), Plérin (France), Rennes (France), Villejuif (France), Augsburg (Germany), Berlin (Germany), Braunschweig (Germany), Dresden (Germany), Dusseldorf (Germany), Emmendingen (Germany), Jena (Germany), Köln (Germany), Lubeck (Germany), Mannheim (Germany), Nürtingen (Germany), Tuebingen (Germany), Wuppertal (Germany), Cork (Ireland), Dublin (Ireland), Haifa (Israel), Jerusalem (Israel), Kfar Saba (Israel), Petach Tikva (Israel), Tel Aviv (Israel), Tel Hashomer (Israel), Arezzo (Italy), Faenza (Italy), Meldola (Italy), Milano (Italy), Modena (Italy), Roma (Italy), Terni (Italy), Trento (Italy), Barcelona (Spain), Córdoba (Spain), Guadalajara (Spain), Lugo (Spain), Madrid (Spain), Málaga (Spain), Oviedo (Spain), Sabadell (Spain), Santander (Spain), Sevilla (Spain), Valencia (Spain), Cardiff (United Kingdom), Taunton (United Kingdom), Wirral (United Kingdom)

NCT03732820
PHASE 3

Study on Olaparib Plus Abiraterone as First-line Therapy in Men With Metastatic Castration-resistant Prostate Cancer

TARGETS
PARP, CYP17

LOCATIONS: Alabama, Alaska, Calgary (Canada), Edmonton (Canada), Arizona, Kelowna (Canada), California, Colorado, Connecticut, Illinois, Indiana, Louisiana, Michigan, Missouri, Montana, Nebraska, New Jersey, New York, North Carolina, Halifax (Canada), London (Canada), Toronto (Canada), Pennsylvania, Greenfield Park (Canada), Montreal (Canada), South Carolina, Texas, Wisconsin, Box Hill (Australia), Darlinghurst (Australia), Greenslopes (Australia), Herston (Australia), Kingswood (Australia), Kurralta Park (Australia), St Albans (Australia), Waratah (Australia), Gent (Belgium), Belo Horizonte (Brazil), Curitiba (Brazil), Fortaleza (Brazil), Porto Alegre (Brazil), Rio de Janeiro (Brazil), Sao Paulo (Brazil), São José do Rio Preto (Brazil), Santiago (Chile), Temuco (Chile), Viña del Mar (Chile), Brno (Czechia), Praha (Czechia), Praha 5 (Czechia), Angers Cedex 01 (France), BESANCON Cedex (France), Caen Cedex 05 (France), Marseille (France), Pierre Benite (France), Quimper Cedex (France), Toulouse Cedex 3 (France), Vandoeuvre les Nancy (France), Bergisch Gladbach (Germany), Bremen (Germany), Duisburg (Germany), Freiburg im Breisgau (Germany), Heinsberg (Germany), Köln (Germany), Mettmann (Germany), Nürnberg (Germany), Nürtingen (Germany), Ulm (Germany), Milano (Italy), Napoli (Italy), Orbassano (Italy), Pavia (Italy), Bunkyo-ku (Japan), Hirakata-shi (Japan), Kanazawa-shi (Japan), Kashiwara-shi (Japan), Kawagoe-shi (Japan), Kita-gun (Japan), Kyoto-shi (Japan), Maebashi-shi (Japan), Miyazaki-city (Japan), Nagoya-shi (Japan), Osaka-shi (Japan), Osakasayama-shi (Japan), Sagami-hara-shi (Japan), Sakura-shi (Japan), Shinjuku-ku (Japan), Toon-shi (Japan), Yokohama-shi (Japan), Daegu (Korea, Republic of), Goyang-si (Korea, Republic of), Seoul (Korea, Republic of), Hilversum (Netherlands), Nijmegen (Netherlands), Tilburg (Netherlands), Bratislava (Slovakia), Presov (Slovakia), Sala (Slovakia), Trencin (Slovakia), Barcelona (Spain), Gerona (Spain), Madrid (Spain), Malaga (Spain), Sevilla (Spain), Adana (Turkey), Ankara (Turkey), Istanbul (Turkey), Izmir (Turkey), Karsiyaka (Turkey), Guildford (United Kingdom), Manchester (United Kingdom), Sheffield (United Kingdom), Southampton (United Kingdom), Swansea (United Kingdom)

NCT02952534
PHASE 2

A Study of Rucaparib in Patients With Metastatic Castration-resistant Prostate Cancer and Homologous Recombination Gene Deficiency

TARGETS
PARP

LOCATIONS: Alabama, Arizona, Slough (United Kingdom), California, Colorado, Connecticut, Delaware, District of Columbia, Northwood (United Kingdom), Florida, Georgia, Illinois, Louisiana, Maryland, Massachusetts, Michigan, Minnesota, Missouri, Nebraska, Nevada, New Jersey, Saint Leonards (Australia), New York, North Carolina, Ohio, Hamilton (Canada), London (Canada), Ottawa (Canada), Oregon, Pennsylvania, Sutton (United Kingdom), Hobart (Australia), Tennessee, Texas, Frankston (Australia), Geelong (Australia), Malvern (Australia), Virginia, Washington, Miranda (Australia), Orange (Australia), Subiaco (Australia), Wagga Wagga (Australia), Antwerp (Belgium), Gent (Belgium), Kortrijk (Belgium), Liège (Belgium), Roeselare (Belgium), Toronto (Canada), Copenhagen (Denmark), Herlev (Denmark), Vejle (Denmark), Caen (France), Dijon (France), Le Mans (France), Lille (France), Nancy (France), Paris (France), Plérin (France), Rennes (France), Augsburg (Germany), Berlin (Germany), Dresden (Germany), Dusseldorf (Germany), Emmendingen (Germany), Hamburg (Germany), Heidelberg (Germany), Jena (Germany), Köln (Germany), Lübeck (Germany), Mannheim (Germany), Nürtingen (Germany), Tuebingen (Germany), Wuppertal (Germany), Cork (Ireland), Dublin (Ireland), Haifa (Israel), Jerusalem (Israel), Kfar Saba (Israel), Petach Tikva (Israel), Ramat Gan (Israel), Tel Aviv (Israel), Arezzo (Italy), Faenza (Italy), Milano (Italy), Modena (Italy), Rome (Italy), Terni (Italy), Trento (Italy), Badalona (Spain), Barcelona (Spain), Guadalajara (Spain), Lugo (Spain), Madrid (Spain), Oviedo (Spain), Sabadell (Spain), Santander (Spain), Sevilla (Spain), Valencia (Spain), Headington (United Kingdom), Liverpool (United Kingdom), London (United Kingdom), Southampton (United Kingdom), Taunton (United Kingdom), Wirral (United Kingdom)

PRF#

CLINICAL TRIALS
NCT03330405
PHASE 2

Javelin Parp Medley: Avelumab Plus Talazoparib In Locally Advanced Or Metastatic Solid Tumors

TARGETS
PD-L1, PARP

LOCATIONS: Edmonton (Canada), Arkansas, California, District of Columbia, Obninsk (Russian Federation), Massachusetts, Minnesota, Sydney (Australia), New York, Ohio, Toronto (Canada), Brisbane (Australia), Texas, Murdoch (Australia), Brussels (Belgium), Bruxelles (Belgium), Charleroi (Belgium), Copenhagen (Denmark), Herlev (Denmark), Budapest (Hungary), Miskolc (Hungary), Pecs (Hungary), Incheon (Korea, Republic of), Seoul (Korea, Republic of), Chelyabinsk (Russian Federation), Moscow (Russian Federation), Omsk (Russian Federation), Yaroslavl (Russian Federation), Leicester (United Kingdom), London (United Kingdom), Newcastle Upon Tyne (United Kingdom)

NCT03742895
PHASE 2

Efficacy and Safety of Olaparib (MK-7339) in Participants With Previously Treated, Homologous Recombination Repair Mutation (HRRm) or Homologous Recombination Deficiency (HRD) Positive Advanced Cancer (MK-7339-002 / LYNK-002)

TARGETS
PARP

LOCATIONS: Nagoya (Japan), Medellin (Colombia), Arizona, Oradea (Romania), Berazategui (Argentina), Ciudad de Buenos Aires (Argentina), California, Chelyabinsk (Russian Federation), Kashiwa (Japan), Comuna Floresti (Romania), Bogota (Colombia), Craiova (Romania), Georgia, Seongnam-si (Korea, Republic of), Guadalajara (Mexico), Istanbul (Turkey), Kentucky, Trujillo (Peru), Pozuelo de Alarcon (Spain), Maryland, Massachusetts, Michigan, Nebraska, New Jersey, Port Macquarie (Australia), New York, Monterrey (Mexico), Oklahoma, Suita (Japan), Pennsylvania, Montreal (Canada), South Dakota, Darlinghurst (Australia), Madero (Mexico), Utah, Washington, Nedlands (Australia), Buenos Aires (Argentina), Quebec (Canada), Barranquilla (Colombia), Cali (Colombia), Monteria (Colombia), Valledupar (Colombia), Copenhagen (Denmark), Herlev (Denmark), Odense (Denmark), Bordeaux (France), Dijon (France), Nice (France), Poitiers (France), Strasbourg (France), Villejuif (France), Guatemala (Guatemala), Quetzaltenango (Guatemala), Cork (Ireland), Dublin (Ireland), Haifa (Israel), Jerusalem (Israel), Ramat Gan (Israel), Tel Aviv (Israel), Napoli (Italy), Rozzano (Italy), Siena (Italy), Kyoto (Japan), Tokyo (Japan), Seoul (Korea, Republic of), Chihuahua (Mexico), Mexico City (Mexico), Oaxaca (Mexico), Santiago De Quetaro (Mexico), Lima (Peru), Brasov (Romania), Bucuresti (Romania), Cluj Napoca (Romania), Arkhangelsk (Russian Federation), Kazan (Russian Federation), Moscow (Russian Federation), Ryazan (Russian Federation), Saint Petersburg (Russian Federation), Saint-Petersburg (Russian Federation), Samara (Russian Federation), St.Petersburg (Russian Federation), Barcelona (Spain), Bellinzona (Switzerland), Geneva (Switzerland), Zuerich (Switzerland), Adana (Turkey), Ankara (Turkey), Antalya (Turkey), Edirne (Turkey), Izmir (Turkey), Konya (Turkey), Manchester (United Kingdom), Newcastle-upon-Tyne (United Kingdom), Oxford (United Kingdom), Sheffield (United Kingdom)

NCT03565991
PHASE 2

Javelin BRCA/ATM: Avelumab Plus Talazoparib in Patients With BRCA or ATM Mutant Solid Tumors

TARGETS
PD-L1, PARP

LOCATIONS: Torette Di Ancona (Italy), California, Kashiwa (Japan), Meldola (Italy), Georgia, Louisiana, Monza (Italy), Milano (Italy), Massachusetts, Missouri, Pamplona (Spain), New Jersey, New York, Amsterdam (Netherlands), Ohio, Oklahoma, Pennsylvania, Tennessee, Texas, Chuo-ku (Japan), Rotterdam (Netherlands), Brussel (Belgium), Brussels (Belgium), Edegem (Belgium), Copenhagen (Denmark), Odense C (Denmark), Clermont Ferrand (France), La Rochelle (France), Montpellier Cedex 5 (France), Napoli (Italy), Roma (Italy), Barcelona (Spain), Madrid (Spain), Sevilla (Spain), London (United Kingdom)

NCT02854436
PHASE 2

An Efficacy and Safety Study of Niraparib in Men With Metastatic Castration-Resistant Prostate Cancer and DNA-Repair Anomalies

TARGETS
PARP

LOCATIONS: Arizona, Vancouver (Canada), California, Colorado, Illinois, Kentucky, Louisiana, Massachusetts, Michigan, New York, North Carolina, Toronto (Canada), Pennsylvania, Montreal (Canada), South Carolina, Texas, Virginia, Washington, Wisconsin, Camperdown (Australia), Darlinghurst (Australia), East Albury (Australia), Kurralta Park (Australia), Melbourne (Australia), Murdoch (Australia), North Ryde (Australia), Port Macquarie (Australia), Randwick (Australia), Wahroonga (Australia), Aalst (Belgium), Brussel (Belgium), Charleroi (Belgium), Gent (Belgium), Haine-Saint-Paul, La Louviere (Belgium), Hasselt (Belgium), Kortrijk (Belgium), Liège (Belgium), Namur (Belgium), Ottignies (Belgium), Wilrijk (Belgium), Barretos (Brazil), Belo Horizonte (Brazil), Curitiba (Brazil), Fortaleza (Brazil), Ijuí (Brazil), Itajaí (Brazil), Joinville (Brazil), Natal (Brazil), Salvador (Brazil), Sao Paulo (Brazil), Quebec (Canada), Avignon Cedex 9 (France), Besancon (France), Caen (France), Lyon (France), Nice Cedex 2 (France), Paris (France), Reims (France), Strasbourg (France), Villejuif Cedex (France), Beer-Sheva (Israel), Haifa (Israel), Kfar Saba (Israel), Ramat Gan (Israel), Zrifin (Israel), Seoul (Korea, Republic of), Amsterdam (Netherlands), Groningen (Netherlands), Maastricht (Netherlands), Rotterdam (Netherlands), Moscow (Russian Federation), Omsk (Russian Federation), Tomsk (Russian Federation), Barcelona (Spain), Córdoba (Spain), Madrid (Spain), Málaga (Spain), Pozuelo de Alarcon (Spain), Santander (Spain), Santiago de Compostela (Spain), Sevilla (Spain), Valencia (Spain), Lund (Sweden), Stockholm (Sweden), Umeå (Sweden), Kaohsiung (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Taipei (Taiwan), Taoyuan County (Taiwan), Blackburn (United Kingdom), Bristol (United Kingdom), Cardiff (United Kingdom), Exeter (United Kingdom), London (United Kingdom), Preston (United Kingdom)

PRF#

CLINICAL TRIALS

NCT03521037
PHASE 1

Rucaparib Hepatic Impairment Study in Patients With a Solid Tumor

TARGETS
PARP
LOCATIONS: Biała Podlaska (Poland), Poznań (Poland), Szczecin (Poland), Warszawa (Poland), Bratislava (Slovakia), Newcastle Upon Tyne (United Kingdom)

PRF#

CLINICAL TRIALS

GENE
RET

ALTERATION
amplification

RATIONALE
RET amplification, activating mutations, or activating fusions may confer sensitivity to kinase

inhibitors targeting RET.

NCT02029001

PHASE 2

Adapting Treatment to the Tumor Molecular Alterations for Patients With Advanced Solid Tumors: My Own Specific Treatment

TARGETS
mTOR, FLT3, KIT, PDGFRs, RAFs, RET, VEGFRs, ERBB2, EGFR, FGFR1, FGFR2, FGFR3, PARP, PD-L1, CTLA-4

LOCATIONS: Bordeaux (France), Lyon (France), Marseille (France), Paris (France), Pierre-Bénite (France), Toulouse (France)

NCT03297606

PHASE 2

Canadian Profiling and Targeted Agent Utilization Trial (CAPTUR)

TARGETS
VEGFRs, ABL, SRC, ALK, AXL, MET, ROS1, TRKA, TRKC, DDR2, KIT, PDGFRs, EGFR, PD-1, CTLA-4, PARP, CDK4, CDK6, CSF1R, FLT3, RET, mTOR, ERBB2, ERBB3, BRAF, MEK, SMO

LOCATIONS: Vancouver (Canada), Kingston (Canada), London (Canada), Ottawa (Canada), Toronto (Canada), Montreal (Canada), Regina (Canada), Saskatoon (Canada)

NCT02616185

PHASE 1

A Phase 1 Study To Evaluate Escalating Doses Of A Vaccine-Based Immunotherapy Regimen For Prostate Cancer (PrCa VBIR)

TARGETS
CSF1R, FLT3, KIT, PDGFRs, RET, VEGFRs, CTLA-4, PD-1

LOCATIONS: Arizona, Connecticut, Nebraska, New Jersey, New York, North Carolina, Pennsylvania, Washington

PRF#

CLINICAL TRIALS

GENE
TMPRSS2

RATIONALE
Fusion of TMPRSS2 with ETS family
transcription factors may indicate sensitivity to

inhibitors of HDAC. TMPRSS2-ERG fusions may
also indicate sensitivity to PARP inhibitors.

ALTERATION
TMPRSS2-ERG fusion

NCT03395197

PHASE 3

Talazoparib + Enzalutamide vs. Enzalutamide Monotherapy in mCRPC (TALAPRO-2)

TARGETS
PARP

LOCATIONS: Santiago de Compostela (Spain), Nagoya (Japan), Alaska, Hiroasaki (Japan), Arizona, Tauranga (New Zealand), L'Hospitalet de Llobregat (Spain), Sabadell (Spain), Pergamino (Argentina), Cremona (Italy), California, Christchurch (New Zealand), Kashiwa (Japan), Colorado, Port Elizabeth (South Africa), Meldola (Italy), Goyang-si (Korea, Republic of), Sapporo (Japan), Indiana, Kanazawa (Japan), Yokohama (Japan), Yokosuka (Japan), Kentucky, Nebraska, New Jersey, New Mexico, Camperdown (Australia), Darlinghurst (Australia), Port Macquarie (Australia), New York, Ohio, Osaka-shi (Japan), Osakasayama (Japan), Suita (Japan), Pennsylvania, Auchenflower (Australia), Brisbane (Australia), Chermide (Australia), South Brisbane (Australia), Southport (Australia), Rosario (Argentina), Hamamatsu (Japan), South Carolina, Terni (Italy), Orbassano (Italy), Tennessee, Texas, Meguro-ku (Japan), Shinjuku-ku (Japan), Utah, Vina del Mar (Chile), Melbourne (Australia), North Melbourne (Australia), Hamilton (New Zealand), Washington, Cape Town (South Africa), George (South Africa), Kraaifontein, Cape Town (South Africa), Wisconsin, Cordoba (Argentina), Gent (Belgium), Kortrijk (Belgium), Yvoir (Belgium), Ostrava-Poruba (Czechia), Helsinki (Finland), Kempele (Finland), Kuopio (Finland), Oulu (Finland), Tampere (Finland), Turku (Finland), Suresnes (France), Suresnes Cedex (France), VILLEJUIF cedex (France), Kirchheim (Germany), Muenster (Germany), Nuertingen (Germany), Ravensburg (Germany), Reutlingen (Germany), Budapest (Hungary), Debrecen (Hungary), Pecs (Hungary), Tel Aviv (Israel), Napoli (Italy), Chiba (Japan), Fukuoka (Japan), Kagoshima (Japan), Kumamoto (Japan), Tokushima (Japan), Yamagata (Japan), Busan (Korea, Republic of), Daegu (Korea, Republic of), Seoul (Korea, Republic of), Auckland (New Zealand), Lorenskog (Norway), Norbyhagen (Norway), Oslo (Norway), Trondheim (Norway), Arequipa (Peru), Lima (Peru), Gdynia (Poland), Otwock (Poland), Warszawa (Poland), Barcelona (Spain), Solna (Sweden), Stockholm (Sweden), Umeå (Sweden), Cornwall (United Kingdom), Cronwall (United Kingdom), Glasgow (United Kingdom), London (United Kingdom)

NCT03748641

PHASE 3

A Study of Niraparib in Combination With Abiraterone Acetate and Prednisone Versus Abiraterone Acetate and Prednisone for Treatment of Participants With Metastatic Prostate Cancer

TARGETS
CYP17, PARP

LOCATIONS: Calgary (Canada), Arizona, Vancouver (Canada), Victoria (Canada), California, Colorado, Connecticut, Florida, Kentucky, Maryland, Massachusetts, Nebraska, Nevada, New Jersey, New York, Hamilton (Canada), Kingston (Canada), Toronto (Canada), Pennsylvania, Montreal (Canada), South Carolina, Texas, Utah, Virginia, Washington, Buenos Aires (Argentina), Capital Federal (Argentina), Ciudad Autonoma Buenos Aires (Argentina), Ciudad Autonoma Buenos Aires (Argentina), Cordoba (Argentina), Mar del Plata (Argentina), Pergamino (Argentina), Rosario (Argentina), San Salvador de Juiuy (Argentina), Adelaide (Australia), Benowa (Australia), Birtinya (Australia), Brisbane (Australia), Darlinghurst (Australia), Hobart (Australia), Macquarie University (Australia), Malvern (Australia), Melbourne (Australia), Murdoch (Australia), Nedlands (Australia), Randwick (Australia), Wahroonga (Australia), Wollongong (Australia), Aalst (Belgium), Antwerpen (Belgium), Charleroi (Belgium), Gent (Belgium), Liege (Belgium), Merksem (Belgium), Natal (Brazil), Rio de Janeiro (Brazil), Sao Paulo (Brazil), Sorocaba (Brazil), São Paulo (Brazil), Pleven (Bulgaria), Plovdiv (Bulgaria), Sofia (Bulgaria), Vratsa (Bulgaria), Hradec Králove (Czechia), Liberec (Czechia), Olomouc (Czechia), Pardubice (Czechia), Plzen (Czechia), Praha 2 (Czechia), Praha 4 (Czechia), Besancon (France), Bordeaux (France), Clermont Ferrand (France), Lyon (France), Nancy (France), Nice Cedex 2 (France), Saint Herblain (France), Saint Mande (France), Strasbourg (France), Braunschweig (Germany), Duisburg (Germany), Magdeburg (Germany), Muenster (Germany), Nuertingen (Germany), Budapest (Hungary), Debrecen (Hungary), Gyula (Hungary), Nyiregyhaza (Hungary), Szeged (Hungary), Beer Yaakov (Israel), Haifa (Israel), Petah Tikva (Israel), Ramat Gan (Israel), Faenza (Italy), Milano (Italy), Napoli (Italy), Padova (Italy), Parma (Italy), Perugia (Italy), Pisa (Italy), Roma (Italy), Terni (Italy), Torino (Italy), Trento (Italy), Busan (Korea, Republic of), Daegu (Korea, Republic of), Daejeon (Korea, Republic of), Goyang-Si (Korea, Republic of), Gwangju (Korea, Republic of), Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Suwon (Korea, Republic of), George Town (Malaysia), Johor Bahru (Malaysia), Kota Kinabalu (Malaysia), Kuala Lumpur (Malaysia), Kuching (Malaysia), Aguascalientes (Mexico), Durango (Mexico), Monterrey (Mexico), Zapopan (Mexico), Amsterdam (Netherlands), Groningen (Netherlands), Leidschendam (Netherlands), Nieuwegein (Netherlands), Nijmegen (Netherlands), Rotterdam (Netherlands), Sittard-Geleen (Netherlands), Bydgoszcz (Poland), Gdańsk (Poland), Gdynia (Poland), Lodz (Poland), Siedlce (Poland), Szczecin (Poland), Warszawa (Poland), Wrocław (Poland), Barnaul (Russian Federation), Ivanovo (Russian Federation), Kursk (Russian Federation), Moscow (Russian Federation), Nizhny Novgorod (Russian Federation), Omsk (Russian Federation), Pyatigorsk (Russian Federation), Saint Petersburg (Russian Federation), Saint-Petersburg (Russian Federation), Saransk (Russian Federation), Sochi (Russian Federation), Tambov (Russian Federation), Tomsk (Russian Federation), Tyumen (Russian Federation), Vologda (Russian Federation), Johannesburg (South Africa), Pretoria (South Africa), Barcelona (Spain), Coruña (Spain), Jerez de la Frontera (Spain), Madrid (Spain), Málaga (Spain), Pozuelo de Alarcon (Spain), Sabadell (Spain), Santander (Spain), Valencia (Spain), Stockholm (Sweden), Uppsala (Sweden), Kaohsiung (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Taipei (Taiwan), Taoyuan (Taiwan), Adana (Turkey), Ankara (Turkey), Antalya (Turkey), Edirne (Turkey), Istanbul (Turkey), Izmir (Turkey), Kocaeli (Turkey), Cherkasy (Ukraine), Dnipro (Ukraine), Dnipro (Ukraine), Ivano-Frankivsk (Ukraine), Khakhiv (Ukraine), Kharkiv (Ukraine), Kyiv (Ukraine), Lviv (Ukraine), Poltava (Ukraine), Uzhgorod (Ukraine), Blackburn (United Kingdom), Lancaster (United Kingdom), London (United Kingdom), Torquay (United Kingdom), Truro (United Kingdom), Wolverhampton (United Kingdom)

PRF#

CLINICAL TRIALS
NCT03732820
PHASE 3

Study on Olaparib Plus Abiraterone as First-line Therapy in Men With Metastatic Castration-resistant Prostate Cancer

TARGETS
PARP, CYP17

LOCATIONS: Alabama, Alaska, Calgary (Canada), Edmonton (Canada), Arizona, Kelowna (Canada), California, Colorado, Connecticut, Illinois, Indiana, Louisiana, Michigan, Missouri, Montana, Nebraska, New Jersey, New York, North Carolina, Halifax (Canada), London (Canada), Toronto (Canada), Pennsylvania, Greenfield Park (Canada), Montreal (Canada), South Carolina, Texas, Wisconsin, Box Hill (Australia), Darlinghurst (Australia), Greenslopes (Australia), Herston (Australia), Kingswood (Australia), Kurralt Park (Australia), St Albans (Australia), Waratah (Australia), Gent (Belgium), Belo Horizonte (Brazil), Curitiba (Brazil), Fortaleza (Brazil), Porto Alegre (Brazil), Rio de Janeiro (Brazil), Sao Paulo (Brazil), São José do Rio Preto (Brazil), Santiago (Chile), Temuco (Chile), Viña del Mar (Chile), Brno (Czechia), Praha (Czechia), Praha 5 (Czechia), Angers Cedex 01 (France), BESANCON Cedex (France), Caen Cedex 05 (France), Marseille (France), Pierre Benite (France), Quimper Cedex (France), Toulouse Cedex 3 (France), Vandoeuvre les Nancy (France), Bergisch Gladbach (Germany), Bremen (Germany), Duisburg (Germany), Freiburg im Breisgau (Germany), Heinsberg (Germany), Köln (Germany), Mettmann (Germany), Nürnberg (Germany), Nürtingen (Germany), Ulm (Germany), Milano (Italy), Napoli (Italy), Orbassano (Italy), Pavia (Italy), Bunkyo-ku (Japan), Hirakata-shi (Japan), Kanazawa-shi (Japan), Kashiwara-shi (Japan), Kawagoe-shi (Japan), Kita-gun (Japan), Kyoto-shi (Japan), Maebashi-shi (Japan), Miyazaki-city (Japan), Nagoya-shi (Japan), Osaka-shi (Japan), Osakasayama-shi (Japan), Sagami-hara-shi (Japan), Sakura-shi (Japan), Shinjuku-ku (Japan), Toon-shi (Japan), Yokohama-shi (Japan), Daegu (Korea, Republic of), Goyang-si (Korea, Republic of), Seoul (Korea, Republic of), Hilversum (Netherlands), Nijmegen (Netherlands), Tilburg (Netherlands), Bratislava (Slovakia), Presov (Slovakia), Sala (Slovakia), Trencin (Slovakia), Barcelona (Spain), Gerona (Spain), Madrid (Spain), Malaga (Spain), Sevilla (Spain), Adana (Turkey), Ankara (Turkey), Istanbul (Turkey), Izmir (Turkey), Karsiyaka (Turkey), Guildford (United Kingdom), Manchester (United Kingdom), Sheffield (United Kingdom), Southampton (United Kingdom), Swansea (United Kingdom)

NCT03330405
PHASE 2

Javelin Parp Medley: Avelumab Plus Talazoparib In Locally Advanced Or Metastatic Solid Tumors

TARGETS
PD-L1, PARP

LOCATIONS: Edmonton (Canada), Arkansas, California, District of Columbia, Obninsk (Russian Federation), Massachusetts, Minnesota, Sydney (Australia), New York, Ohio, Toronto (Canada), Brisbane (Australia), Texas, Murdoch (Australia), Brussels (Belgium), Bruxelles (Belgium), Charleroi (Belgium), Copenhagen (Denmark), Herlev (Denmark), Budapest (Hungary), Miskolc (Hungary), Pecs (Hungary), Incheon (Korea, Republic of), Seoul (Korea, Republic of), Chelyabinsk (Russian Federation), Moscow (Russian Federation), Omsk (Russian Federation), Yaroslavl (Russian Federation), Leicester (United Kingdom), London (United Kingdom), Newcastle Upon Tyne (United Kingdom)

NCT03521037
PHASE 1

Rucaparib Hepatic Impairment Study in Patients With a Solid Tumor

TARGETS
PARP

LOCATIONS: Biąła Podlaska (Poland), Poznań (Poland), Szczecin (Poland), Warszawa (Poland), Bratislava (Slovakia), Newcastle Upon Tyne (United Kingdom)

PRF#

CLINICAL TRIALS
NCT03834519
PHASE 3

Study of Pembrolizumab (MK-3475) Plus Olaparib Versus Abiraterone Acetate or Enzalutamide in Metastatic Castration-resistant Prostate Cancer (mCRPC) (MK-7339-010/KEYLYNK-010)

TARGETS
PD-1, CYP17, PARP, AR

LOCATIONS: Edmonton (Canada), L Hospitalet De Llobregat (Spain), Kelowna (Canada), Vancouver (Canada), Berazategui (Argentina), Buenos Aires (Argentina), California, Cambridge (United Kingdom), Avignon (France), Kashiwa (Japan), Sakura (Japan), Matsuyama (Japan), Caceres (Spain), Georgia, Illinois, Kanazawa (Japan), Hwasun Gun (Korea, Republic of), Sagami-hara (Japan), Yokohama (Japan), Louisiana, Pozuelo de Alarcon (Spain), Maryland, Massachusetts, Michigan, Rozzano (Italy), Montana, Kashi-hara (Japan), Nebraska, Nevada, New Mexico, Darlinghurst (Australia), Macquarie University (Australia), Port Macquarie (Australia), Waratah (Australia), Wollongong (Australia), North Carolina, Halifax (Canada), Ohio, Hamilton (Canada), Toronto (Canada), Osakasayama (Japan), Suita (Japan), Rimouski (Canada), Sherbrooke (Canada), Herston (Australia), Tugun (Australia), Ijuí (Brazil), São Paulo (Brazil), Hidaka (Japan), Koshigaya (Japan), Itajai (Brazil), Rosario (Argentina), São José do Rio Preto (Brazil), Hamamatsu (Japan), South Carolina, Utah, Box Hill (Australia), Melbourne (Australia), Virginia, Murdoch (Australia), Wisconsin, Ube (Japan), Caba (Argentina), Córdoba (Argentina), Graz (Austria), Linz (Austria), Salzburg (Austria), Wien (Austria), Quebec (Canada), Santiago (Chile), Temuco (Chile), Amiens (France), Besançon (France), Bordeaux (France), Brest (France), Clermont-Ferrand (France), Lyon (France), Marseille (France), Montpellier (France), Nancy (France), Orleans (France), Paris (France), Pierre Benite (France), Saint Quentin (France), Suresnes (France), Toulouse (France), Villejuif (France), Berlin (Germany), Duesseldorf (Germany), Erlangen (Germany), Muenchen (Germany), Nuernberg (Germany), Nuertingen (Germany), Trier (Germany), Tuebingen (Germany), Dublin (Ireland), Limerick (Ireland), Be'er- Ya'akov (Israel), Haifa (Israel), Jerusalem (Israel), Kfar Saba (Israel), Petach-Tikwa (Israel), Ramat Gan (Israel), Tel-Aviv (Israel), Bologna (Italy), Catania (Italy), Roma (Italy), Terni (Italy), Trento (Italy), Chiba (Japan), Fukuoka (Japan), Miyazaki (Japan), Nagasaki (Japan), Tokyo (Japan), Goyang-si (Korea, Republic of), Seoul (Korea, Republic of), Almelo (Netherlands), Amsterdam (Netherlands), Hoofddorp (Netherlands), Leeuwarden (Netherlands), Leidschendam (Netherlands), Nijmegen (Netherlands), Rotterdam (Netherlands), Auckland (New Zealand), Chelyabinsk (Russian Federation), Krasnoyarsk (Russian Federation), Moscow (Russian Federation), Omsk (Russian Federation), Saint Petersburg (Russian Federation), Samara (Russian Federation), Tomsk (Russian Federation), Barcelona (Spain), Girona (Spain), Malaga (Spain), Sabadell (Spain), Taichung (Taiwan), Tainan (Taiwan), Taipei (Taiwan), Bristol (United Kingdom), Sutton (United Kingdom)

NCT03148795
PHASE 2

A Study of Talazoparib in Men With DNA Repair Defects and Metastatic Castration-Resistant Prostate Cancer

TARGETS
PARP

LOCATIONS: Arizona, California, Suresnes (France), Besançon Cedex (France), Meldola (Italy), Georgia, Seongnam-si (Korea, Republic of), Pozuelo de Alarcon (Spain), Northwood (United Kingdom), Missouri, Pamplona (Spain), Darlinghurst (Australia), Northmead (Australia), Westmead (Australia), North Carolina, Auchenflower (Australia), Cairns (Australia), Chermiside (Australia), South Brisbane (Australia), Southport (Australia), Ijuí (Brazil), Porto Alegre (Brazil), Roma (Italy), Barretos (Brazil), Jau (Brazil), South Carolina, Sutton (United Kingdom), Nijmegen (Netherlands), Texas, Orbassano (Italy), Linz (Austria), Mestre (Italy), Box Hill (Australia), Clayton (Australia), Frankston (Australia), Heidelberg (Australia), Virginia, Washington, Wisconsin, Salzburg (Austria), Vienna (Austria), Bruxelles (Belgium), Gent (Belgium), Leuven (Belgium), Angers Cedex 02 (France), Angers Cedex 2 (France), Angers cedex 02 (France), Bordeaux cedex (France), La Roche sur Yon (France), Le Mans Cedex 02 (France), Strasbourg (France), Strasbourg Cedex (France), Villejuif Cedex (France), Mannheim (Germany), Nuertingen (Germany), Tubingen (Germany), Wuerzburg (Germany), Budapest (Hungary), Debrecen (Hungary), Nyiregyhaza (Hungary), Cremona (Italy), Naples (Italy), Padova (Italy), Parma (Italy), Torino (Italy), Busan (Korea, Republic of), Daegu (Korea, Republic of), Seoul (Korea, Republic of), Brzozow (Poland), Kielce (Poland), Barcelona (Spain), Malaga (Spain), Valencia (Spain), Cambridge (United Kingdom)

NCT03431350
PHASE 1/2

A Study of Niraparib Combination Therapies for the Treatment of Metastatic Castration-Resistant Prostate Cancer

TARGETS
PARP, PD-1, CYP17

LOCATIONS: Calgary (Canada), Arizona, Vancouver (Canada), Colorado, Indiana, Maryland, Michigan, New York, Toronto (Canada), Pennsylvania, Montreal (Canada), South Carolina, Tennessee, Texas, Utah, Virginia, Wisconsin, Aalst (Belgium), Antwerpen (Belgium), Brussels (Belgium), Gent (Belgium), Kortrijk (Belgium), Liege (Belgium), Beer Yaakov (Israel), Beer-Sheva (Israel), Haifa (Israel), Petach Tikva (Israel), Ramat Gan (Israel), Firenze (Italy), Lecce (Italy), Macerata (Italy), Milano (Italy), Napoli (Italy), Barcelona (Spain), Madrid (Spain), Malaga (Spain), Bath (United Kingdom), London (United Kingdom), Southampton (United Kingdom), Sutton (United Kingdom), Truro (United Kingdom)

NCT02264678
PHASE 1/2

Ascending Doses of AZD6738 in Combination With Chemotherapy and/or Novel Anti Cancer Agents

TARGETS
ATR, PARP, PD-L1

LOCATIONS: California, New York, Saint Herblain (France), Villejuif (France), Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Cambridge (United Kingdom), London (United Kingdom), Sutton (United Kingdom), Withington (United Kingdom)

PRF#

CLINICAL TRIALS

NCT03338790

PHASE 2

An Investigational Immunotherapy Study of Nivolumab in Combination With Rucaparib, Docetaxel, or Enzalutamide in Metastatic Castration-resistant Prostate Cancer

TARGETS
PD-1, PARP, AR

LOCATIONS: Kelowna (Canada), Caba (Argentina), Capital Federal (Argentina), Ciudad Autonoma de Buenos Aires (Argentina), Monteria (Colombia), Leon (Mexico), Guadalajara (Mexico), Santiago de Chile (Chile), Moncton (Canada), Camperdown (Australia), Westmead (Australia), Hamilton (Canada), South Brisbane (Australia), Porto Alegre (Brazil), Culiacan (Mexico), Elizabeth Vale (Australia), Vina Del Mar (Chile), Clayton (Australia), Heidelberg (Australia), Cordoba (Argentina), Montreal (Canada), Quebec (Canada), Medellin (Colombia), Besancon (France), Clermont-ferrand (France), Lyon (France), Marseille (France), Villejuif (France)

PRF#

APPENDIX

Variants of Unknown Significance

NOTE One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

BRCA2
N863S and amplification

IKBKE
N700S

KDR
A1065T

MAF
G145S

NBN
amplification

PALB2
V78I

The content provided as a professional service by Foundation Medicine, Inc., has not been reviewed or approved by the FDA.

Electronically signed by Douglas Lin, M.D. | Julia Elvin, M.D., Ph.D., Laboratory Director Foundation Medicine, Inc. | Roche Customer Care: +49 7624 14 2098 or europe.foundationmedicine@roche.com

Sample Preparation: FMI Germany GmbH, Nonnenwald 2, 82377 Penzberg, Germany
Sample Analysis: FMI Germany GmbH, Nonnenwald 2, 82377 Penzberg, Germany

PRF#

APPENDIX

Genes Assayed in FoundationOne®CDx

FoundationOne CDx is designed to include genes known to be somatically altered in human solid tumors that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay interrogates 324 genes as well as introns of 36 genes involved in rearrangements. The assay will be updated periodically to reflect new knowledge about cancer biology.

DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS

ABL1	ACVR1B	AKT1	AKT2	AKT3	ALK	ALOX12B	AMER1 (FAM123B)	APC
AR	ARAF	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX	AURKA
AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2	BCL6
BCOR	BCORL1	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2
BTB	C11orf30 (EMSY)	C17orf39 (GID4)	CALR	CARD11	CASP8	CBFB	CBL	CCND1
CCND2	CCND3	CCNE1	CD22	CD274 (PD-L1)	CD70	CD79A	CD79B	CDC73
CDH1	CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B
CDKN2C	CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R
CTCF	CTNNA1	CTNNB1	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1
DDR2	DIS3	DNMT3A	DOT1L	EED	EGFR	EP300	EPHA3	EPHB1
EPHB4	ERBB2	ERBB3	ERBB4	ERCC4	ERG	ERRF1	ESR1	EZH2
FAM46C	FANCA	FANCC	FANCG	FANCL	FAS	FBXW7	FGF10	FGF12
FGF14	FGF19	FGF23	FGF3	FGF4	FGF6	FGFR1	FGFR2	FGFR3
FGFR4	FH	FLCN	FLT1	FLT3	FOXO2	FUBP1	GABRA6	GATA3
GATA4	GATA6	GNA11	GNA13	GNAQ	GNAS	GRM3	GSK3B	H3F3A
HDAC1	HGF	HNF1A	HRAS	HSD3B1	ID3	IDH1	IDH2	IGF1R
IKBKE	IKZF1	INPP4B	IRF2	IRF4	IRS2	JAK1	JAK2	JAK3
JUN	KDMSA	KDMS5C	KDM6A	KDR	KEAP1	KEL	KIT	KLHL6
KMT2A (MLL)	KMT2D (MLL2)	KRAS	LTK	LYN	MAF	MAP2K1 (MEK1)	MAP2K2 (MEK2)	MAP2K4
MAP3K1	MAP3K13	MAPK1	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1
MERTK	MET	MITF	MKNK1	MLH1	MPL	MRE11A	MSH2	MSH3
MSH6	MST1R	MTAP	MTOR	MUTYH	MYC	MYCL (MYCL1)	MYCN	MYD88
NBN	NF1	NF2	NFE2L2	NFKB1A	NKX2-1	NOTCH1	NOTCH2	NOTCH3
NPM1	NRAS	NSD3 (WHSC1L1)	NT5C2	NTRK1	NTRK2	NTRK3	P2RY8	PALB2
PARK2	PARP1	PARP2	PARP3	PAX5	PBRM1	PDCD1 (PD-1)	PDCD1LG2 (PD-L2)	PDGFRA
PDGFRB	PDK1	PIK3C2B	PIK3C2G	PIK3CA	PIK3CB	PIK3R1	PIM1	PMS2
POLD1	POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI	PTCH1
PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51	RAD51B	RAD51C
RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10	REL	RET
RICTOR	RNF43	ROS1	RPTOR	SDHA	SDHB	SDHC	SDHD	SETD2
SF3B1	SGK1	SMAD2	SMAD4	SMARCA4	SMARCB1	SMO	SNCAIP	SOC3
SOX2	SOX9	SPEN	SPOP	SRC	STAG2	STAT3	STK11	SUFU
SYK	TBX3	TEK	TET2	TGFBR2	TIPARP	TNFAIP3	TNFRSF14	TP53
TSC1	TSC2	TYRO3	U2AF1	VEGFA	VHL	WHSC1	WT1	XPO1
XRCC2	ZNF217	ZNF703						

DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS

ALK	BCL2	BCR	BRAF	BRCA1	BRCA2	CD74	EGFR	ETV4
ETV5	ETV6	EWSR1	EZR	FGFR1	FGFR2	FGFR3	KIT	KMT2A (MLL)
MSH2	MYB	MYC	NOTCH2	NTRK1	NTRK2	NUTM1	PDGFRA	RAF1
RARA	RET	ROS1	RSP02	SDC4	SLC34A2	TERC*	TERT**	TPR2SS2

*TERC is an NCRNA

**Promoter region of TERT is interrogated

ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER GENOMIC SIGNATURES

Loss of Heterozygosity (LOH) score

Microsatellite (MS) status

Tumor Mutational Burden (TMB)

FoundationOne CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Cipalstraat 3, 2440 Geel, Belgium.



ABOUT FOUNDATIONONE CDx

FoundationOne CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details:
www.rochefoundationmedicine.com/f1cdxtech.

INTENDED USE

FoundationOne®CDx (F1CDx) is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 324 genes and select gene rearrangements, as well as genomic signatures including microsatellite instability (MSI), tumor mutational burden (TMB), and for selected forms of ovarian cancer, loss of heterozygosity (LOH) score, using DNA isolated from formalin-fixed, paraffin-embedded (FFPE) tumor tissue specimens. The test is intended as a companion diagnostic to identify patients who may benefit from treatment with therapies in accordance with approved therapeutic product labeling. Additionally, F1CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with solid malignant neoplasms.

TEST PRINCIPLES

FoundationOne CDx will be performed exclusively as a laboratory service using DNA extracted from formalin-fixed, paraffin-embedded (FFPE) tumor samples. The proposed assay will employ a single DNA extraction method from routine FFPE biopsy or surgical resection specimens, 50-1000 ng of which will undergo whole-genome shotgun library construction and hybridization-based capture of all coding exons from 309 cancer-related genes, one promoter region, one non-coding (ncRNA), and select intronic regions from 34 commonly rearranged genes, 21 of which also include the coding exons. The assay therefore includes detection of alterations in a total of 324 genes. Using an Illumina® HiSeq platform, hybrid

capture-selected libraries will be sequenced to high uniform depth (targeting >500X median coverage with >99% of exons at coverage >100X). Sequence data will be processed using a customized analysis pipeline designed to accurately detect all classes of genomic alterations, including base substitutions, indels, focal copy number amplifications, homozygous gene deletions, and selected genomic rearrangements (e.g., gene fusions). Additionally, genomic signatures including loss of heterozygosity (LOH), microsatellite instability (MSI) and tumor mutational burden (TMB) will be reported.

THE REPORT

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. The F1CDx report may be used as an aid to inform molecular eligibility for clinical trials. Note: The association of a therapy with a genomic alteration or signature does not necessarily indicate pharmacologic effectiveness (or lack thereof); no association of a therapy with a genomic alteration or signature does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness).

Diagnostic Significance

FoundationOne CDx identifies alterations to select cancer-associated genes or portions of genes (biomarkers). In some cases, the Report also highlights selected negative test results regarding biomarkers of clinical significance.

Qualified Alteration Calls (Equivocal and Subclonal)

An alteration denoted as "amplification - equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne CDx for identifying a copy number amplification is four (4) for ERBB2 and six (6) for all other genes. Conversely, an alteration denoted as "loss - equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as "subclonal" is one that the FoundationOne CDx analytical methodology has identified as being present in <10% of the assayed tumor DNA.

Ranking of Alterations and Therapies

Genomic Signatures and Gene Alterations
Therapies are ranked based on the following

criteria: Therapies approved in the EU in patient's tumor type (ranked alphabetically within each NCCN category) followed by therapies approved in the EU in another tumor type (ranked alphabetically within each NCCN category).

Clinical Trials

Pediatric trial qualification → Geographical proximity → Later trial phase.

NCCN Categorization

Genomic signatures and gene alterations detected may be associated with certain National Comprehensive Cancer Network (NCCN) Compendium drugs or biologics (www.nccn.org). The NCCN categories indicated reflect the highest possible category for a given therapy in association with each genomic signature or gene alteration. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories please refer to the NCCN Compendium.

Limitations

1. The MSI-H/MSS designation by FMI F1CDx test is based on genome wide analysis of 95 microsatellite loci and not based on the 5 or 7 MSI loci described in current clinical practice guidelines. The threshold for MSI-H/MSS was determined by analytical concordance to comparator assays (IHC and PCR) using uterine, cecum and colorectal cancer FFPE tissue. The clinical validity of the qualitative MSI designation has not been established. For Microsatellite Instability (MSI) results, confirmatory testing using a validated orthogonal method should be considered.
2. TMB by F1CDx is defined based on counting the total number of all synonymous and nonsynonymous variants present at 5% allele frequency or greater (after filtering) and reported as mutations per megabase (mut/Mb) unit rounded to the nearest integer. The clinical validity of TMB defined by this panel has not been established.
3. The LOH score is determined by analyzing SNPs spaced at 1Mb intervals across the genome on the FoundationOne CDx test and extrapolating an LOH profile, excluding arm- and chromosome-wide LOH segments. Detection of LOH has been verified only for ovarian cancer patients, and the LOH score result may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas. The LOH score will be reported as "Cannot Be Determined" if the sample is not of sufficient quality to confidently determine LOH.

PRF#

APPENDIX

About FoundationOne®CDx

Performance of the LOH classification has not been established for samples below 35% tumor content. There may be potential interference of ethanol with LOH detection. The interfering effects of xylene, hemoglobin, and triglycerides on the LOH score have not been demonstrated.

Sample

LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

NO GUARANTEE OF CLINICAL BENEFIT

This Report makes no promises or guarantees that a particular drug will be effective in the treatment of disease in any patient. This Report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne CDx.

TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report. Certain sample or variant characteristics may result in reduced sensitivity. FoundationOne CDx is performed using DNA derived from tumor, and as such germline events may not be reported.

SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
mut/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

PDF Service version: 2.6.0

The median exon coverage for this sample is 1,003x

APPENDIX

References

PRF#

1. Gatalica Z, Snyder C, Maney T, et al. ePub Dec 2014 (2014) PMID: 25392179
2. Kroemer G, Galluzzi L, Zitvogel L, et al. 4 (7):e1058597 (2015) PMID: 26140250
3. Lal N, Beggs AD, Willcox BE, et al. 4 (3):e976052 (2015) PMID: 25949894
4. Le DT, Uram JN, Wang H, et al. ePub Jun 2015 (2015) PMID: 26028255
5. ASCO-SITC 2016; Abstract P60
6. Hall et al., 2016; ASCO Abstract 1523
7. Cheng et al., 2016; ASCO Abstract 251
8. Pritchard CC, Morrissey C, Kumar A, et al. ePub Sep 2014 (2014) PMID: 25255306
9. Schwarzenbach H, Chun FK, Müller I, et al. ePub Jul 2008 (2008) PMID: 18336598
10. Suzuki H, Komiya A, Aida S, et al. 86 (10):956-61 (1995) PMID: 7493915
11. Dominguez-Valentin M, Joost P, Therkildsen C, et al. ePub Mar 2016 (2016) PMID: 27013479
12. Rosty C, Walsh MD, Lindor NM, et al. ePub Dec 2014 (2014) PMID: 25117503
13. Ahman AK, Jonsson BA, Damber JE, et al. 87 (4):334-8 (2001) PMID: 11251526
14. Kocarnik JM, Shiovitz S, Phipps AI 3 (4):269-76 (2015) PMID: 26337942
15. You JF, Buhard O, Ligtenberg MJ, et al. ePub Dec 2010 (2010) PMID: 21081928
16. Bairwa NK, Saha A, Gochhait S, et al. ePub 2014 (2014) PMID: 24623249
17. Boland CR, Thibodeau SN, Hamilton SR, et al. 58 (22):5248-57 (1998) PMID: 9823339
18. Pawlik TM, Raut CP, Rodriguez-Bigas MA 20 (4-5):199-206 (2004) PMID: 15528785
19. Boland CR, Goel A ePub Jun 2010 (2010) PMID: 20420947
20. null ePub 12 2017 (2017) PMID: 29101156
21. Hellmann MD, Callahan MK, Awad MM, et al. ePub May 2018 (2018) PMID: 29731394
22. Hellmann MD, Ciuleanu TE, Pluzanski A, et al. ePub 05 2018 (2018) PMID: 29658845
23. Hellmann MD, Nathanson T, Rizvi H, et al. ePub May 2018 (2018) PMID: 29657128
24. Ready N, Hellmann MD, Awad MM, et al. ePub Apr 2019 (2019) PMID: 30785829
25. Goodman AM, Kato S, Bazhenova L, et al. ePub 11 2017 (2017) PMID: 28835386
26. Carbone DP, Reck M, Paz-Ares L, et al. ePub 06 2017 (2017) PMID: 28636851
27. Rizvi H, Sanchez-Vega F, La K, et al. ePub Mar 2018 (2018) PMID: 29337640
28. Forde PM, Chaff JE, Smith KN, et al. ePub 05 2018 (2018) PMID: 29658848
29. Samstein RM, Lee CH, Shoushtari AN, et al. ePub 02 2019 (2019) PMID: 30643254
30. Miao D, Margolis CA, Vokes NI, et al. ePub Sep 2018 (2018) PMID: 30150660
31. Chae YK, Davis AA, Raparia K, et al. ePub 03 2019 (2019) PMID: 30425022
32. Colli LM, Machiela MJ, Myers TA, et al. ePub 07 2016 (2016) PMID: 27197178
33. Wang VE, Urisman A, Albacker L, et al. ePub 09 2017 (2017) PMID: 28923100
34. Gandara DR, Paul SM, Kowanetz M, et al. ePub 09 2018 (2018) PMID: 30082870
35. Rosenberg JE, Hoffman-Censits J, Powles T, et al. ePub May 2016 (2016) PMID: 26952546
36. Balar AV, Galsky MD, Rosenberg JE, et al. ePub 01 2017 (2017) PMID: 27939400
37. Powles T, Durán I, van der Heijden MS, et al. ePub Feb 2018 (2018) PMID: 29268948
38. Mariathasan S, Turley SJ, Nickles D, et al. ePub 02 2018 (2018) PMID: 29443960
39. Johnson DB, Frampton GM, Rioth MJ, et al. ePub Nov 2016 (2016) PMID: 27671167
40. Eroglu Z, Zaretsky JM, Hu-Lieskován S, et al. ePub 01 2018 (2018) PMID: 29320474
41. Shin DS, Zaretsky JM, Escuin-Ordinas H, et al. ePub 02 2017 (2017) PMID: 27903500
42. Riaz N, Havel JJ, Makarov V, et al. ePub Nov 2017 (2017) PMID: 29033130
43. Cristescu R, Mogg R, Ayers M, et al. ePub 10 2018 (2018) PMID: 30309915
44. Fabrizio DA, George TJ, Dunne RF, et al. 9 (4):610-617 (2018) PMID: 30151257
45. Hanna GJ, Lizotte P, Cavanaugh M, et al. ePub Feb 2018 (2018) PMID: 29467336
46. Ott PA, Bang YJ, Piha-Paul SA, et al. ePub Feb 2019 (2019) PMID: 30557521
47. Mehner JM, Panda A, Zhong H, et al. ePub Jun 2016 (2016) PMID: 27159395
48. Santin AD, Bellone S, Buza N, et al. 22 (23):5682-5687 (2016) PMID: 27486176
49. Snyder A, Makarov V, Merghoub T, et al. ePub Dec 2014 (2014) PMID: 25409260
50. Van Allen EM, Miao D, Schilling B, et al. ePub Oct 2015 (2015) PMID: 25959337
51. Roszik J, Haydu LE, Hess KR, et al. ePub 10 2016 (2016) PMID: 27776519
52. Chalmers ZR, Connelly CF, Fabrizio D, et al. ePub 04 2017 (2017) PMID: 28420421
53. Lawrence MS, Stojanov P, Polak P, et al. ePub Jul 2013 (2013) PMID: 23770567
54. Alexandrov LB, Nik-Zainal S, Wedge DC, et al. ePub Aug 2013 (2013) PMID: 23945592
55. Barbieri CE, Baca SC, Lawrence MS, et al. ePub May 2012 (2012) PMID: 22610119
56. Baca SC, Prandi D, Lawrence MS, et al. ePub Apr 2013 (2013) PMID: 23622249
57. null ePub Nov 2015 (2015) PMID: 26544944
58. Grasso CS, Wu YM, Robinson DR, et al. ePub Jul 2012 (2012) PMID: 2272839
59. Gundem G, Van Loo P, Kremeyer B, et al. ePub Apr 2015 (2015) PMID: 25830880
60. Robinson D, Van Allen EM, Wu YM, et al. ePub May 2015 (2015) PMID: 26000489
61. Pfeifer GP, You YH, Besaratinia A 571 (1-2):19-31 (2005) PMID: 15748635
62. Hill VK, Gartner JJ, Samuels Y, et al. ePub 2013 (2013) PMID: 23875803
63. Pfeifer GP, Denissenko MF, Olivier M, et al. 21 (48):7435-51 (2002) PMID: 12379884
64. Rizvi NA, Hellmann MD, Snyder A, et al. ePub Apr 2015 (2015) PMID: 25765070
65. Cancer Genome Atlas Research Network, Kandoth C, Schultz N, et al. ePub May 2013 (2013) PMID: 23636398
66. Briggs S, Tomlinson I ePub Jun 2013 (2013) PMID: 23447401
67. Heitzer E, Tomlinson I ePub Feb 2014 (2014) PMID: 24583393
68. null ePub Jul 2012 (2012) PMID: 22810696
69. Roberts SA, Gordenin DA ePub 12 2014 (2014) PMID: 25568919
70. Patel JC, Maughan BL, Agarwal AM, et al. 2013 :981684 (2013) PMID: 23819055
71. Chowdhury S, Kirby R 257 (1760):15-8, 2 (2013) PMID: 23724747
72. Ahmed SR, Brooman PJ, Shalet SM, et al. 2 (8347):415-9 (1983) PMID: 6135909
73. Smith MR, Saad F, Chowdhury S, et al. ePub Apr 2018 (2018) PMID: 29420164
74. Fizazi K, Shore N, Tammela TL, et al. ePub 03 2019 (2019) PMID: 30763142
75. Fizazi K, Massard C, Bono P, et al. ePub Aug 2014 (2014) PMID: 24974051
76. Soifer HS, Souleimanian N, Wu S, et al. ePub Feb 2012 (2012) PMID: 22174412
77. null ePub Aug 2011 (2011) PMID: 21808270
78. van der Steen T, Tindall DJ, Huang H 14 (7):14833-59 (2013) PMID: 23863692
79. Lee E, Madar A, David G, et al. ePub Sep 2013 (2013) PMID: 24019458
80. Gowda PS, Deng JD, Mishra S, et al. ePub Nov 2013 (2013) PMID: 23989930
81. Liu LL, Xie N, Sun S, et al. ePub Jun 2014 (2014) PMID: 23851510
82. Wadosky KM, Koochekpour S ePub Sep 2016 (2016) PMID: 27487144
83. Bartek J, Mistrik M, Bartkova J ePub Nov 2013 (2013) PMID: 24203954
84. van Soest RJ, van Royen ME, de Morree ES, et al. ePub Dec 2013 (2013) PMID: 24200698
85. Scher HI, Lu D, Schreiber NA, et al. ePub Nov 2016 (2016) PMID: 27262168
86. Antonarakis ES, Lu C, Lubner B, et al. ePub Aug 2015 (2015) PMID: 26181238
87. Loddick SA, Ross SJ, Thomason AG, et al. ePub Sep 2013 (2013) PMID: 23861347
88. Rathkopf DE, Morris MJ, Fox JJ, et al. ePub Oct 2013 (2013) PMID: 24002508
89. Smith MR, Antonarakis ES, Ryan CJ, et al. ePub 12 2016 (2016) PMID: 27160947
90. Rathkopf DE, Antonarakis ES, Shore ND, et al. 23 (14):3544-3551 (2017) PMID: 28213364
91. Soh SF, Huang CK, Lee SO, et al. ePub Jan 2014 (2014) PMID: 24042123
92. Bianchini D, Omlin A, Pezaro C, et al. ePub Nov 2013 (2013) PMID: 24169353
93. Li H, Hassona MD, Lack NA, et al. ePub Nov 2013 (2013) PMID: 23939374
94. McKay RR, Werner L, Mostaghel EA, et al. 23 (4):935-945 (2017) PMID: 27683182
95. Asangani IA, Dommeti VL, Wang X, et al. ePub Jun 2014 (2014) PMID: 24759320
96. Asangani IA, Wilder-Romans K, Dommeti VL, et al. ePub 04 2016 (2016) PMID: 26792867
97. Faviere EJ, Wilcox D, Lin X, et al. ePub 01 2017 (2017) PMID: 27707886
98. Chan SC, Selth LA, Li Y, et al. ePub Jul 2015 (2015) PMID: 25908785
99. Piha-Paul SA, Sachdev JC, Barve M, et al. (2019) PMID: 31420359
100. Montgomery B, Eisenberger MA, Rettig MB, et al. 22 (6):1356-63 (2016) PMID: 26527750
101. Palmberg C, Koivisto P, Kakkola L, et al. 164 (6):1992-5 (2000) PMID: 11061898
102. Visakorpi T, Hyytinen E, Koivisto P, et al. 9 (4):401-6 (1995) PMID: 7795646
103. Koivisto P, Kononen J, Palmberg C, et al. 57 (2):314-9 (1997) PMID: 9000575

104. Haapala K, Kuukasjärvi T, Hyytinen E, et al. 38 (3):474-8 (2007) PMID: 17217995
105. Schrecengost R, Knudsen KE ePub Jun 2013 (2013) PMID: 23806491
106. Schmidt LJ, Tindall DJ ePub Apr 2013 (2013) PMID: 23565753
107. Taylor BS, Schultz N, Hieronymus H, et al. ePub Jul 2010 (2010) PMID: 20579941
108. Koivisto PA, Helin HJ 189 (2):219-23 (1999) PMID: 10547578
109. Niraula S, Chi K, Joshua AM ePub Apr 2012 (2012) PMID: 22124843
110. Kaufman B, Shapira-Frommer R, Schmutzler RK, et al. ePub Jan 2015 (2015) PMID: 25366685
111. Mateo J, Carreira S, Sandhu S, et al. ePub Oct 2015 (2015) PMID: 26510020
112. Tutt A, Robson M, Garber JE, et al. ePub Jul 2010 (2010) PMID: 20609467
113. Robson M, Im SA, Senkus E, et al. ePub 08 2017 (2017) PMID: 28578601
114. Mirza MR, Monk BJ, Herrstedt J, et al. ePub 12 2016 (2016) PMID: 27717299
115. Sandhu SK, Schelman WR, Wilding G, et al. ePub Aug 2013 (2013) PMID: 23810788
116. Swisher EM, Lin KK, Oza AM, et al. ePub Jan 2017 (2017) PMID: 27908594
117. Drew Y, Ledermann J, Hall G, et al. ePub Mar 2016 (2016) PMID: 27002934
118. Pujade-Lauraine E, Ledermann JA, Selle F, et al. ePub Sep 2017 (2017) PMID: 28754483
119. Ledermann JA, Harter P, Gourley C, et al. ePub Nov 2016 (2016) PMID: 27617661
120. Fong PC, Boss DS, Yap TA, et al. ePub Jul 2009 (2009) PMID: 19553641
121. Audeh MW, Carmichael J, Penson RT, et al. ePub Jul 2010 (2010) PMID: 20609468
122. Fong PC, Yap TA, Boss DS, et al. ePub May 2010 (2010) PMID: 20406929
123. Gelmon KA, Tischkowitz M, Mackay H, et al. ePub Sep 2011 (2011) PMID: 21862407
124. Kaye SB, Lubinski J, Matulonis U, et al. ePub Feb 2012 (2012) PMID: 22203755
125. Domchek SM, Aghajanian C, Shapira-Frommer R, et al. ePub Feb 2016 (2016) PMID: 26723501
126. Moore K, Colombo N, Scambia G, et al. ePub Oct 2018 (2018) PMID: 30345884
127. Konstantinopoulos et al., 2018; ASCO Abstract 106
128. Kummar et al., 2019; AACR Abstract CT138/4
129. Isakoff SJ, Mayer EL, He L, et al. ePub Jun 2015 (2015) PMID: 25847936
130. Dann RB, DeLoia JA, Timms KM, et al. ePub Jun 2012 (2012) PMID: 22406760
131. Evers B, Drost R, Schut E, et al. 14 (12):3916-25 (2008) PMID: 18559613
132. Chung JH, Dewal N, Sokol E, et al. 3 (2019) PMID: 31218271
133. Pritchard CC, Mateo J, Walsh MF, et al. ePub Aug 2016 (2016) PMID: 27433846
134. Mitra AV, Bancroft EK, Barbachano Y, et al. ePub Jan 2011 (2011) PMID: 20840664
135. Castro E, Goh C, Olmos D, et al. ePub May 2013 (2013) PMID: 23569316
136. Castro E, Goh C, Leongamornlert D, et al. ePub Aug 2015 (2015) PMID: 25454609
137. Castro E, Romero-Laorden N, Del Pozo A, et al. ePub Feb 2019 (2019) PMID: 30625039
138. Cheng HH, Pritchard CC, Boyd T, et al. ePub 06 2016 (2016) PMID: 26724258
139. null ePub 06 2016 (2016) PMID: 26708042
140. Yang H, Jeffrey PD, Miller J, et al. ePub Sep 2002 (2002) PMID: 12228710
141. null ePub Jul 2011 (2011) PMID: 21731065
142. Al Abo M, Dejsuphong D, Hirota K, et al. ePub Feb 2014 (2014) PMID: 24285729
143. Biswas K, Das R, Alter BP, et al. ePub Sep 2011 (2011) PMID: 21719596
144. Biswas K, Das R, Egginton JM, et al. ePub Sep 2012 (2012) PMID: 22678057
145. Claes K, Poppe B, Coene I, et al. 90 (6):1244-51 (2004) PMID: 15026808
146. Cote S, Arcand SL, Royer R, et al. ePub Jan 2012 (2012) PMID: 21947752
147. Fackenthal JD, Cartegni L, Krainer AR, et al. 71 (3):625-31 (2002) PMID: 12145750
148. Farrugia DJ, Agarwal MK, Pankratz VS, et al. ePub May 2008 (2008) PMID: 18451181
149. Gómez García EB, Oosterwijk JC, Timmermans M, et al. ePub 2009 (2009) PMID: 19200354
150. Guidugli L, Pankratz VS, Singh N, et al. ePub Jan 2013 (2013) PMID: 23108138
151. Kluska A, Balabas A, Paziewska A, et al. ePub May 2015 (2015) PMID: 25948282
152. Kuznetsov SG, Liu P, Sharan SK ePub Aug 2008 (2008) PMID: 18607349
153. Roy R, Chun J, Powell SN ePub Dec 2011 (2011) PMID: 22193408
154. Siaud N, Barbera MA, Egashira A, et al. ePub Dec 2011 (2011) PMID: 22194698
155. Thomassen M, Kruse TA, Jensen PK, et al. 10 (2):116-20 (2006) PMID: 16792514
156. Xia B, Sheng Q, Nakanishi K, et al. 22 (6):719-29 (2006) PMID: 16793542
157. Guidugli L, Shimelis H, Masica DL, et al. ePub 02 2018 (2018) PMID: 29394989
158. Fraile-Bethencourt E, Díez-Gómez B, Velásquez-Zapata V, et al. ePub Mar 2017 (2017) PMID: 28339459
159. Miki Y, Swensen J, Shattuck-Eidens D, et al. 266 (5182):66-71 (1994) PMID: 7545954
160. Wooster R, Bignell G, Lancaster J, et al. 378 (6559):789-92 (null) PMID: 8524414
161. Ford D, Easton DF, Bishop DT, et al. 343 (8899):692-5 (1994) PMID: 7907678
162. null ePub Jun 2005 (2005) PMID: 16369438
163. Struwing JP, Hartge P, Wacholder S, et al. 336 (20):1401-8 (1997) PMID: 9145676
164. Bougie O, Weberpals JI ePub 2011 (2011) PMID: 22312502
165. null 91 (15):1310-6 (1999) PMID: 10433620
166. Hahn SA, Greenhalf B, Ellis I, et al. 95 (3):214-21 (2003) PMID: 12569143
167. Monnerat C, Chompret A, Kannengiesser C, et al. 6 (4):453-61 (2007) PMID: 17624602
168. Casula M, Muggiano A, Cossu A, et al. ePub Oct 2009 (2009) PMID: 19799798
169. Moran A, O'Hara C, Khan S, et al. ePub Jun 2012 (2012) PMID: 22187320
170. Whittmore AS, Gong G, Itnyre J 60 (3):496-504 (1997) PMID: 9042908
171. Claus EB, Schildkraut JM, Thompson WD, et al. 77 (11):2318-24 (1996) PMID: 8635102
172. Oddoux C, Struwing JP, Clayton CM, et al. 14 (2):188-90 (1996) PMID: 8841192
173. King MC, Marks JH, Mandell JB, et al. ePub Oct 2003 (2003) PMID: 14576434
174. Hall MJ, Reid JE, Burbidge LA, et al. 115 (10):2222-33 (2009) PMID: 19241424
175. Lin JJ, Kennedy E, Sequist LV, et al. ePub 11 2016 (2016) PMID: 27544060
176. Drilon A, Rekhman N, Arcila M, et al. ePub Dec 2016 (2016) PMID: 27825636
177. Drilon A, Wang L, Hasanovic A, et al. ePub Jun 2013 (2013) PMID: 23533264
178. Drilon A, Wang L, Arcila ME, et al. 21 (16):3631-9 (2015) PMID: 25567908
179. Elisei R, Schlumberger MJ, Müller SP, et al. ePub Oct 2013 (2013) PMID: 24002501
180. Gautschi O, Milia J, Filleron T, et al. ePub May 2017 (2017) PMID: 28447912
181. Mukhopadhyay S, Pennell NA, Ali SM, et al. ePub Nov 2014 (2014) PMID: 25436805
182. Michels S, Scheel AH, Scheffler M, et al. ePub Jan 2016 (2016) PMID: 26762747
183. Sherman SI, Clary DO, Elisei R, et al. ePub Dec 2016 (2016) PMID: 27525386
184. Schlumberger M, Jarzab B, Cabanillas ME, et al. 22 (1):44-53 (2016) PMID: 26311725
185. Jones SJ, Laskin J, Li YY, et al. ePub 2010 (2010) PMID: 20696054
186. Wu H, Shih JY, Yang JC ePub Sep 2015 (2015) PMID: 26291023
187. null 5 (4):7-11 (2015) PMID: 26693082
188. Falchook GS, Ordóñez NG, Bastida CC, et al. ePub 05 2016 (2016) PMID: 25366691
189. Gautschi O, Zander T, Keller FA, et al. ePub May 2013 (2013) PMID: 23584301
190. Lee SH, Lee JK, Ahn MJ, et al. ePub 02 2017 (2017) PMID: 27803005
191. Subbiah V, Berry J, Roxas M, et al. ePub Jul 2015 (2015) PMID: 25982012
192. Wells SA, Robinson BG, Gagel RF, et al. ePub Jan 2012 (2012) PMID: 22025146
193. Yoh K, Seto T, Satouchi M, et al. ePub 01 2017 (2017) PMID: 27825616
194. Subbiah V, Meric-Bernstam F, Mills GB, et al. ePub Aug 2014 (2014) PMID: 25085632
195. Dawson DM, Lawrence EG, MacLennan GT, et al. 90 (7):519-23 (1998) PMID: 9539247
196. Takahashi M, Ritz J, Cooper GM 42 (2):581-8 (1985) PMID: 2992805
197. Gao J, Aksoy BA, Dogrusoz U, et al. ePub Apr 2013 (2013) PMID: 23550210
198. Björkman M, Iljin K, Halonen P, et al. ePub Dec 2008 (2008) PMID: 18798265
199. Brenner JC, Ateeq B, Li Y, et al. ePub May 2011 (2011) PMID: 21575865
200. Chatterjee P, Choudhary GS, Sharma A, et al. ePub 2013 (2013) PMID: 23565244
201. Hussain M, Carducci MA, Slovin S, et al. ePub Oct 2014 (2014) PMID: 24764124
202. Carver BS, Tran J, Gopalan A, et al. ePub May 2009 (2009) PMID: 19396168
203. King JC, Xu J, Wongvipat J, et al. ePub May 2009 (2009) PMID: 19396167
204. Scheble VJ, Braun M, Beroukhim R, et al. ePub Aug 2010 (2010) PMID: 20473283
205. Wang JJ, Liu YX, Wang W, et al. ePub 2012 (2012) PMID: 23244085
206. Tomlins SA, Rhodes DR, Perner S, et al. ePub Oct 2005 (2005) PMID: 16254181

207. Sun QP, Li LY, Chen Z, et al. ePub Sep 2010 (2010) PMID: 20616363
208. Lara et al., 2016; ASCO Abstract 5037
209. Wang J, Cai Y, Ren C, et al. ePub Sep 2006 (2006) PMID: 16951141
210. Kissick HT, Sanda MG, Dunn LK, et al. ePub Dec 2013 (2013) PMID: 24149465
211. Qu X, Randhawa G, Friedman C, et al. ePub 2013 (2013) PMID: 24098661
212. Johnson H, Zhou M, Osunkoya AO ePub Oct 2013 (2013) PMID: 23849895
213. Tavukcu HH, Mangir N, Ozyurek M, et al. 54 (6):359-63 (2013) PMID: 23789042
214. Cheng L, Davidson DD, MacLennan GT, et al. ePub Oct 2013 (2013) PMID: 23887154
215. Green WM, Hicks JL, De Marzo A, et al. ePub Sep 2013 (2013) PMID: 23664537
216. Clark JP, Cooper CS ePub Aug 2009 (2009) PMID: 19657377
217. Reig Ó, Marín-Aguilera M, Carrera G, et al. ePub 11 2016 (2016) PMID: 26948395
218. Vaarala MH, Porvari KS, Kellokumpu S, et al. 193 (1):134-40 (2001) PMID: 11169526
219. Clinckemalie L, Spans L, Dubois V, et al. ePub Dec 2013 (2013) PMID: 24109594
220. Wang J, Cai Y, Yu W, et al. ePub Oct 2008 (2008) PMID: 18922926
221. Bronner CE, Baker SM, Morrison PT, et al. 368 (6468):258-61 (1994) PMID: 8145827
222. Gazzoli I, Loda M, Garber J, et al. 62 (14):3925-8 (2002) PMID: 12124320
223. Lipson EJ, Sharfman WH, Drake CG, et al. 19 (2):462-8 (2003) PMID: 23169436
224. Chen Y, Wang J, Fraig MM, et al. 22 (5):1033-43 (2003) PMID: 12684669
225. Wilczak W, Rashed S, Hube-Magg C, et al. ePub 01 2017 (2017) PMID: 27803051
226. Kadyrov FA, Dzantiev L, Constantin N, et al. 126 (2):297-308 (2006) PMID: 16873062
227. Tomer G, Buermeyer AB, Nguyen MM, et al. 277 (24):21801-9 (2002) PMID: 11897781
228. Plotz G, Raedle J, Brieger A, et al. ePub Jun 2003 (2003) PMID: 12799449
229. Wu H, Zeng H, Lam R, et al. ePub Aug 2015 (2015) PMID: 26249686
230. Guerrette S, Acharya S, Fishel R 274 (10):6336-41 (1999) PMID: 10037723
231. Lynch HT, Lynch PM, Lanspa SJ, et al. ePub Jul 2009 (2009) PMID: 19659756
232. Raevaara TE, Korhonen MK, Lohi H, et al. 129 (2):537-49 (2005) PMID: 16083711
233. Silva FC, Valentin MD, Ferreira Fde O, et al. ePub Jan 2009 (2009) PMID: 19466295
234. Sehgal R, Sheahan K, O'Connell PR, et al. 5 (3):497-507 (2014) PMID: 24978665
235. null 4 (3):233-7 (2005) PMID: 16136383
236. Zehir A, Benayed R, Shah RH, et al. ePub Jun 2017 (2017) PMID: 28481359
237. Yan M, Xu H, Waddell N, et al. ePub Apr 2012 (2012) PMID: 22537934
238. Xu H, Yan M, Patra J, et al. ePub Jan 2011 (2011) PMID: 21255398
239. Stevens KN, Wang X, Fredericksen Z, et al. ePub Sep 2011 (2011) PMID: 21607584
240. Sehl ME, Langer LR, Papp JC, et al. 15 (6):2192-203 (2009) PMID: 19276285
241. Supernat A, Lapińska-Szumczyk S, Sawicki S, et al. 4 (4):727-732 (2012) PMID: 23205091
242. Deb S, Xu H, Tuynman J, et al. ePub Mar 2014 (2014) PMID: 24548858
243. Supernat A, Lapińska-Szumczyk S, Majewska H, et al. 7 (5):613-9 (2014) PMID: 25048628
244. Porkka KP, Tammela TL, Vessella RL, et al. 39 (1):1-10 (2004) PMID: 14603436
245. Davis SJ, Sheppard KE, Anglesio MS, et al. ePub Jun 2015 (2015) PMID: 25852062
246. Atienza JM, Roth RB, Rosette C, et al. 4 (3):361-8 (2005) PMID: 15767545
247. Xu H, Tomaszewski JM, McKay MJ ePub 03 2011 (2011) PMID: 21326324
248. Hill VK, Kim JS, Waldman T 1866 (1):1-11 (2016) PMID: 27207471
249. Solomon DA, Kim JS, Waldman T ePub Jun 2014 (2014) PMID: 24856830
250. Bauerschmidt C, Arrichiello C, Burdak-Rothkamm S, et al. ePub Jan 2010 (2010) PMID: 19906707
251. Yun J, Song SH, Kang JY, et al. ePub Jan 2016 (2016) PMID: 26420833
252. Gelot C, Guirouilh-Barbat J, Lopez BS ePub Jul 2016 (2016) PMID: 27326661
253. Sofueva S, Yaffe E, Chan WC, et al. ePub Dec 2013 (2013) PMID: 24185899
254. Deng Z, Wang Z, Stong N, et al. ePub Nov 2012 (2012) PMID: 23010778
255. Yun J, Song SH, Kim HP, et al. ePub 09 2016 (2016) PMID: 27466323
256. Mahmood SF, Gruel N, Chapeaublanc E, et al. ePub Mar 2014 (2014) PMID: 24148822
257. Hirai H, Arai T, Okada M, et al. ePub Apr 2010 (2010) PMID: 2107315
258. Bridges KA, Hirai H, Buser CA, et al. 17 (17):5638-48 (2011) PMID: 21799033
259. Rajeshkumar NV, De Oliveira E, Ottenhof N, et al. 17 (9):2799-806 (2011) PMID: 21389100
260. Osman AA, Monroe MM, Ortega Alves MV, et al. ePub Feb 2015 (2015) PMID: 25504633
261. Xu L, Huang CC, Huang W, et al. 1 (5):337-46 (2002) PMID: 12489850
262. Xu L, Tang WH, Huang CC, et al. 7 (10):723-34 (2001) PMID: 11713371
263. Camp ER, Wang C, Little EC, et al. ePub Apr 2013 (2013) PMID: 23470564
264. Kim SS, Rait A, Kim E, et al. ePub Feb 2015 (2015) PMID: 25240597
265. Pirolo KF, Nemunaitis J, Leung PK, et al. ePub Sep 2016 (2016) PMID: 27357628
266. Hajdenberg et al., 2012; ASCO Abstract e15010
267. Lehmann S, Bykov VJ, Ali D, et al. ePub Oct 2012 (2012) PMID: 22965953
268. Mohell N, Alfredsson J, Fransson Å, et al. ePub Jun 2015 (2015) PMID: 26086967
269. Fransson Å, Glaessgen D, Alfredsson J, et al. ePub May 2016 (2016) PMID: 27179933
270. Gourley et al., 2016; ASCO Abstract 5571
271. Leijen S, van Geel RM, Pavlick AC, et al. ePub Dec 2016 (2016) PMID: 27601554
272. Moore et al., 2019; ASCO Abstract 5513
273. Leijen S, van Geel RM, Sonke GS, et al. ePub 12 2016 (2016) PMID: 27998224
274. Oza et al., 2015; ASCO Abstract 5506
275. Méndez E, Rodriguez CP, Kao MC, et al. 24 (12):2740-2748 (2018) PMID: 29535125
276. Ma CX, Cai S, Li S, et al. ePub Apr 2012 (2012) PMID: 22446188
277. Kumar A, White TA, MacKenzie AP, et al. ePub Oct 2011 (2011) PMID: 21949389
278. Beltran H, Yelensky R, Frampton GM, et al. ePub May 2013 (2013) PMID: 22981675
279. Schlomm T, Iwers L, Kirstein P, et al. ePub Nov 2008 (2008) PMID: 18552821
280. Liu W, Xie CC, Thomas CY, et al. ePub Jul 2013 (2013) PMID: 23609948
281. Brown CJ, Lain S, Verma CS, et al. ePub Dec 2009 (2009) PMID: 19935675
282. Joerger AC, Fersht AR 77 :557-82 (2008) PMID: 18410249
283. Kato S, Han SY, Liu W, et al. 100 (14):8424-9 (2003) PMID: 12826609
284. Kamada R, Nomura T, Anderson CW, et al. ePub Jan 2011 (2011) PMID: 20978130
285. Bougeard G, Renaux-Petel M, Flaman JM, et al. ePub Jul 2015 (2015) PMID: 26014290
286. Sorrell AD, Espenschied CR, Culver JO, et al. ePub Feb 2013 (2013) PMID: 23355100
287. Nichols KE, Malkin D, Garber JE, et al. 10 (2):83-7 (2001) PMID: 11219776
288. Taubert H, Meyer A, Würfl P 4 (6):365-72 (1998) PMID: 10780879
289. Kleihues P, Schauble B, zur Hausen A, et al. 150 (1):1-13 (1997) PMID: 9006316
290. Gonzalez KD, Noltner KA, Buzin CH, et al. ePub Mar 2009 (2009) PMID: 19204208
291. Laloo F, Varley J, Ellis D, et al. 361 (9363):1101-2 (2003) PMID: 12672316
292. Fizazi K, Tran N, Fein L, et al. ePub Apr 2019 (2019) PMID: 30987939
293. Fizazi K, Tran N, Fein L, et al. ePub 07 2017 (2017) PMID: 28578607
294. Ryan CJ, Smith MR, Fizazi K, et al. ePub Feb 2015 (2015) PMID: 25601341
295. de Bono JS, Logothetis CJ, Molina A, et al. ePub May 2011 (2011) PMID: 21612468
296. Ryan et al., 2016; ASCO Abstract 5061
297. Tagawa ST, Posadas EM, Bruce J, et al. ePub 11 2016 (2016) PMID: 26852075
298. Massard C, Mateo J, Lortet Y, et al. ePub 01 2017 (2017) PMID: 28039155
299. de Wit R, de Bono J, Sternberg CN, et al. ePub Sep 2019 (2019) PMID: 31566937
300. Gucaip A, Tolaney S, Isakoff SJ, et al. 19 (19):5505-12 (2013) PMID: 23965901
301. Akaza H, Hinotsu S, Usami M, et al. 115 (15):3437-45 (2009) PMID: 19536889
302. Bruchovsky N, Klotz L, Crook J, et al. 6 (1):46-52 (2008) PMID: 18501083
303. Calais da Silva F, Calais da Silva FM, Gonçalves F, et al. ePub Aug 2014 (2014) PMID: 23582949
304. Crawford ED, Tombal B, Miller K, et al. ePub Sep 2011 (2011) PMID: 21788033
305. Dawson NA, Figg WD, Cooper MR, et al. 15 (4):1470-7 (1997) PMID: 9193342
306. Horwitz EM, Bae K, Hanks GE, et al. ePub May 2008 (2008) PMID: 18413638
307. Marberger M, Kaisary AV, Shore ND, et al. ePub Apr 2010 (2010) PMID: 20435244
308. Lundström EA, Rencken RK, van Wyk JH, et al. 29 (12):757-65 (2009) PMID: 19888782
309. Smith et al., 2019; ASCO GU Abstract 202

PRF#

APPENDIX

References

310. Del Conte G, Sessa C, von Moos R, et al. ePub Aug 2014 (2014) PMID: 25025963
311. Mateo et al., 2019; ASCO Abstract 5005
312. Antonarakis et al., 2019; ASCO Abstract 5045
313. Clarke N, Wiechno P, Alekseev B, et al. ePub Jul 2018 (2018) PMID: 29880291
314. Karzai F, VanderWeele D, Madan RA, et al. ePub Dec 2018 (2018) PMID: 30514390
315. Yu et al., 2019; ASCO GU Abstract 145
316. Kristeleit et al., 2014; ASCO Abstract 2573
317. Domcheck et al., 2016; ASCO Abstract 4110
318. Abida et al., 2018; ESMO Abstract 793PD
319. Turner et al., 2017; ASCO Abstract 1007
320. Litton JK, Rugo HS, Ettl J, et al. ePub Aug 2018 (2018) PMID: 30110579
321. Ettl J, Quek RGW, Lee KH, et al. ePub Sep 2018 (2018) PMID: 30124753
322. Meehan et al., 2017; AACR Abstract 4687
323. de Bono J, Ramanathan RK, Mina L, et al. ePub 06 2017 (2017) PMID: 28242752
324. Lu E, Thomas GV, Chen Y, et al. ePub 08 2018 (2018) PMID: 30099369